

Coarse differentiation and quasi-isometries of a class of solvable Lie groups II

Irine Peng

Abstract

In this paper, we continue with the results in [P] and compute the group of quasi-isometries for a subclass of split solvable unimodular Lie groups. Consequently, we show that any finitely generated group quasi-isometric to a member of the subclass has to be polycyclic, and is virtually a lattice in an abelian-by-abelian solvable Lie group. We also give an example of a unimodular solvable Lie group that is not quasi-isometric to any finitely generated group, as well deduce some quasi-isometric rigidity results.

Contents

1	Introduction	2
1.1	Proof outline	2
2	Preliminaries	3
2.1	Geometry of a certain class of solvable Lie groups	3
2.2	Notations	5
2.2.1	General remarks about neighborhoods	5
2.2.2	Notations used in split abelian-by-abelian groups	5
3	Shadows, slabs and coarsening	7
3.1	Definitions	8
3.2	Improving almost product map	10
3.3	The constant part of f	14
4	Aligning the linear part of standard maps	17
4.1	S graph, $\hat{S}$ graph and the H graph	20
4.2	Some geometric lemmas	23
4.3	Proof of Theorem 4.1	25
5	Patching	31
5.1	A weak version of an affine map	31
5.2	Consequence of weak height preservation - flats go to flats	34
5.3	Consequences of flats go to flats	36

1 Introduction

A (κ, C) *quasi-isometry* f between metric spaces X and Y is a map $f : X \rightarrow Y$ satisfying

$$\frac{1}{\kappa}d(p, q) - C \leq d(f(p), f(q)) \leq \kappa d(p, q) + C$$

with the additional property that there is a number D such that Y is the D neighborhood of $f(X)$. Two quasi-isometries f, g are considered to be equivalent if there is a number $E > 0$ such that $d(f(p), g(p)) \leq E$ for all $p \in X$.

Let $G = \mathbb{R}^m \rtimes_{\varphi} \mathbb{R}^n$, $G' = \mathbb{R}^{m'} \rtimes_{\varphi'} \mathbb{R}^{n'}$ be connected, simply connected non-degenerate unimodular split solvable groups (See section 2.1 for definitions). We say a map from G to G' is *standard*, if it splits as a product map that respects φ and φ' (See definition 2.1.1). Also homomorphisms φ is called *diagonalizable* if its image can be conjugated into the set of diagonal matrices. The main result of this paper is the following statement.

Theorem 5.2 (abridged) Let G, G' be non-degenerate, unimodular, split abelian-by-abelian solvable Lie groups, and $\phi : G \rightarrow G'$ a κ, C quasi-isometry. Then ϕ is bounded distance from a composition of a left translation and a standard map.

Consequently, *Corollary 5.3.1 If φ is diagonalizable and φ' isn't, then there is no quasi-isometry between them.*

Corollary 1.0.1.

$$\mathcal{QI}(G) = \left(\prod_{[\alpha]} \text{Bilip}(V_{[\alpha]}) \right) \rtimes \text{Sym}(G)$$

Here $[\alpha]$ is an equivalence class of roots and $\text{Sym}(G)$ is a finite group, analogous to the Weyl group in reductive Lie groups, that reflects the symmetries of G . (See section 2.1)

When φ is diagonalizable, as an application the work by Dymarz [D] on quasi-conformal maps on the boundary of G , we have

Corollary 1.0.2. *In the case that φ is diagonalizable, if Γ is a finitely generated group quasi-isometric to a G , then Γ is virtually polycyclic.*

1.1 Proof outline

Our starting point is Theorem 3.1, and our first task is to show that the $\mathbf{A}'$ part of the standard maps (See Definition 2.1.1) f_i are affine. This is done in Section 3, where we will see that the linear part is a scalar multiple of a finite order element in $O(n)$ (where n is the rank of G). We also give interpretations of the linear and constant part of f_i in terms of properties of G' and measure of certain sets in the box where f_i was partially defined. In Section 4, we show that the linear part of the f_i 's in different boxes have to be the same up to scalar multiple in the case that the rank of G is 2 or higher. The rank 1 case is the same as the content in [EFW2]. The proof for higher rank case basically consists of many rank 1 argument as appeared in [EFW2]. In the last section, we put all theses partially defined standard maps together to produce a splitting of the original quasi-isometry.

Acknowledgement I would like to thank Alex Eskin and David Fisher for their incredible patience and assistance. I also wish to thank Tullia Dymarz for helpful conversations and coordinating results of her paper [D] with this one, and to Mikhail Ershov for finding me the reference containing the example of a solvable Lie group that admits no lattices.

2 Preliminaries

Here we recall the settings from [P] and define new terms that will be used in this paper.

2.1 Geometry of a certain class of solvable Lie groups

Non-degenerate, split abelian-by-abelian solvable Lie groups Let $\mathfrak{g}$ be a (real) solvable Lie algebra, and $\mathfrak{a}$ be a Cartan subalgebra. Then there are finitely many non-zero linear functionals $\alpha_i : \mathfrak{a} \rightarrow \mathbb{C}$ called *roots*, such that

$$\mathfrak{g} = \mathfrak{a} \oplus \bigoplus_{\alpha_i} \mathfrak{g}_{\alpha_i}$$

where $\mathfrak{g}_{\alpha_i} = \{x \in \mathfrak{g} : \forall t \in \mathfrak{a}, \exists n, \text{ such that } (ad(t) - \alpha_i(t)Id)^n(x) = 0\}$, Id is the identity map on $\mathfrak{g}$, and $ad : \mathfrak{g} \rightarrow Der_{\mathbb{R}}(\mathfrak{g})$ is the adjoint representation. Let Δ denotes for the set of roots. Then $Aut(\mathfrak{a})$ acts on Δ in a natural way. We define $Perm(\mathfrak{g})$ to be the subgroup consisting $A \in Aut(\mathfrak{a})$ such that

- (i) it leaves the set of roots invariant, i.e. $A\Delta = \Delta$.
- (ii) for every $\alpha \in \Delta$, $\dim \mathfrak{g}_{\alpha \circ A} = \dim \mathfrak{g}_{\alpha}$.

In this way, elements of $Perm(\mathfrak{g})$ induces a permutation on the set Δ , and we define $Sym(\mathfrak{g})$ to be the image of $Perm(\mathfrak{g})$ in the group of permutations of Δ . For a generic $\mathfrak{g}$, its $Perm(\mathfrak{g})$ is trivial.

We say $\mathfrak{g}$ is *split abelian-by-abelian* if $\mathfrak{g}$ is a semidirect product of $\mathfrak{a}$ and $\bigoplus_i \mathfrak{g}_{\alpha_i}$, and both are abelian Lie algebras; *unimodular* if the the roots sum up to zero; and *non-degenerate* if the roots span $\mathfrak{a}^*$. In particular, non-degenerate means that each α_i is real-valued, and the number of roots is at least the dimension of $\mathfrak{a}$. Being unimodular is the same as saying that for every $t \in \mathfrak{a}$, the trace of $ad(t)$ is zero. We extend these definitions to a Lie group if its Lie algebra has these properties, and write $Perm(G)$, $Sym(G)$ to mean $Perm(\mathfrak{g})$ and $Sym(\mathfrak{g})$ where $\mathfrak{g}$ is the Lie algebra of G .

Therefore a connected, simply connected solvable Lie group G that is non-degenerate, split abelian-by-abelian necessary takes the form $G = \mathbf{H} \rtimes_{\varphi} \mathbf{A}$ such that

- (i) both $\mathbf{A}$ and $\mathbf{H}$ are abelian Lie groups.
- (ii) $\varphi : \mathbf{A} \rightarrow Aut(\mathbf{H})$ is injective
- (iii) there are finitely many $\alpha_i \in \mathbf{A}^* \setminus \{0\}$ which together span $\mathbf{A}^*$, and a decomposition of $\mathbf{H} = \bigoplus_i V_{\alpha_i}$
- (iv) there is a basis $\mathcal{B}$ of $\mathbf{H}$ whose intersection with each of V_{α_i} constitute a basis of V_{α_i} , such that for each $\mathbf{t} \in \mathbf{A}$, $\varphi(\mathbf{t})$ with respect to $\mathcal{B}$ is a matrix consists of blocks, one for each V_{α_i} , of the form $e^{\alpha_i(\mathbf{t})}N(\alpha_i(\mathbf{t}))$, where $N(\alpha_i(\mathbf{t}))$ is an upper triangular with 1's on the diagonal and whose off-diagonal entries are polynomials of $\alpha_i(\mathbf{t})$. If in addition, G is unimodular, then $\varphi(\mathbf{t})$ has determinant 1 for all $\mathbf{t} \in \mathbf{A}$.

The *rank* of a non-degenerate, split abelian-by-abelian group G is defined to be the dimension of $\mathbf{A}$, and by a result of Cornulier [C], if two such groups are quasi-isometric, then they have the same rank.

By abuse of notation, we call Δ roots of G as well and coordinatize its points as $((\mathbf{x}_\alpha)_{\alpha \in \Delta}, \mathbf{t})$, where $\mathbf{x}_\alpha = (x_{1,\alpha}, x_{2,\alpha}, \dots, x_{\dim(V_\alpha),\alpha}) \in V_\alpha$, $\mathbf{t} \in \mathbf{A}$. A left invariant Finsler metric that is quasi-isometric to a left invariant Riemannian is given by:

$$d\mathbf{t} + \sum_{\alpha \in \Delta} e^{-\alpha(\mathbf{t})} \left(d\mathbf{x}_\alpha + \sum_j P_{j,\alpha}(\alpha(\mathbf{t})) dx_{j,\alpha} \right)$$

where $P_{j,\alpha}$ is a polynomial. The following consequence is immediate.

Lemma 2.1.1. *If G is non-degenerate, split abelian-by-abelian, then it can be QI embedded into $\prod_{\alpha \in \Delta} H_{\dim(V_\alpha)+1}$, where $H_{s+1} = \mathbb{R}^s \rtimes_\psi \mathbb{R}$ is a non-unimodular solvable Lie group determined by $\psi(t) = e^t N(t)$, where $N(t)$ is a nilpotent matrix (upper triangular with 1's on the diagonal) with polynomial entries, equipped with a left-invariant Finsler metric given by*

$$dt + e^{-t} \left(d\mathbf{x} + \sum_j P_j(t) dx_j \right)$$

where $P_j(t)$ is a polynomial.

Remark 2.1.1. *When $\psi(t)$ is diagonal, H_{s+1} is just the usual hyperbolic space.*

To understand the geometry of H_{s+1} better, we first note that the metric is bilipchitz to one given by $dt + e^{-t}(1 + \max_j P_j(t))d\mathbf{x}$, which is quasi-isometric to one given by $dt + e^{-t}Q(t)d\mathbf{x}$ for some polynomial $Q(t)$. So a function q.i. to the metric on H_{s+1} is the following

$$d((x_1, t_1), (x_2, t_2)) = \begin{cases} |t_1 - t_2| & \text{if } e^{-t_i}Q(t_i)|x_1 - x_2| \leq 1 \text{ for some } i = 1, 2; \\ U_Q(|x_1 - x_2|) - (t_1 + t_2) & \text{otherwise} \end{cases} \quad (1)$$

where $U_Q(|x_1 - x_2|) = t_0$ satisfies

$$e^{-t_0}Q(t_0)|x_1 - x_2| = 1$$

Since exponential grows faster than any polynomials, the function U_Q has the following property:

$$\ln(x) - C_Q \leq U_Q(x) \leq 2 \ln(x) + C_Q \quad (2)$$

for some constant C depends only on the polynomial Q .

Back to the description of G , we declare two roots equivalent if they are positive multiples of each other, and write $[\Xi]$ for the equivalence class containing $\Xi \in \Delta$. Moreover, for $\Xi_i, \Xi_2 \in [\Xi]$, we say Ξ_1 less than Ξ_2 if $\Xi_2/\Xi_1 > 1$. This makes sense because all roots in a root class are positive multiples of each other. A left translate of $V_{[\Xi]} = \oplus_{\sigma \in [\Xi]} V_\sigma$ will be called a *horocycle of root class* $[\Xi]$.

A left translate of $\mathbf{H}$, or a subset of it, is called a *flat*. For $p = (x_\alpha)_\alpha, q = (y_\alpha)_\alpha$ points in $\mathbf{H}$, we compute subsets of $p\mathbf{H}$ and $q\mathbf{H}$ that are within distance 1 of each other according to the embedded metric in Lemma 2.1.1, as the p and q translate of the subset

$$\bigcap_{\alpha \in \Delta: \ln(|x_\alpha - y_\alpha|) \geq 1} \alpha^{-1}[U_\alpha(|x_\alpha - y_\alpha|), \infty] \subset \mathbf{A}$$

Since the roots sum up to zero in a non-degenerate, unimodular, split abelian-by-abelian group, the set where two flats come together can be empty, i.e. the two flats have no intersection. If it is not empty, then the equation above says that it is an unbounded convex subset of $\mathbf{A}$ bounded by hyperplanes parallel to root kernels.

Definition 2.1.1. *Let G, G' be non-degenerate, split abelian-by-abelian Lie groups. A map from G to G' or a subset of them, is called standard map if it takes the form $f \times g$, where $g : \mathbf{H} \rightarrow \mathbf{H}'$ sends foliation by root class horocycles of G to that of G' , and $f : \mathbf{A} \rightarrow \mathbf{A}$ sends foliations by root kernels of G to that of G' . We will often refer to f as the $\mathbf{A}'$ part of a standard map.*

Remark 2.1.2. *Note that when G has at least $\text{rank}(G) + 1$ many root kernels, the condition on f means that f is affine, and when G is rank 1, the condition on f is empty.*

2.2 Notations

2.2.1 General remarks about neighborhoods

Neighborhoods of a set We write $B(p, r)$ for the ball centered at p of radius r , and $N_c(A)$ for the c neighborhood of the set A . We also write $d_H(A, B)$ for the Hausdorff distance between two sets A and B . If $\Omega \subset \mathbb{R}^k$ is a bounded compact set, and $r \in \mathbb{R}$, we write $r\Omega$ for the bounded compact set that is scaled from Ω with respect to the barycenter of Ω .

Given a set X , a point $x_0 \in X$, the (η, C) linear neighborhood of X with respect to x_0 is the set $\{y, s.t. \exists \hat{x} \in X, d(y, \hat{x}) = d(y, X) \leq \eta d(\hat{x}, x_0) + C\}$. Equivalently it is the set $\bigcup_{x \in X} B(x, \eta d(x, x_0) + C)$. By (η, C) linear neighborhood of a set X , we mean the (η, C) linear neighborhood of X with respect to some $x_0 \in X$.

If a quasi-geodesic λ is within (η, C) linear (or just η -linear) neighborhood of a geodesic segment γ , where $\eta \ll 1$ and $C \ll \eta|\lambda|$, then we say that λ admits a geodesic approximation by γ .

2.2.2 Notations used in split abelian-by-abelian groups

Let $G = \mathbf{H} \rtimes \mathbf{A}$ stands for a non-degenerate, unimodular, split abelian-by-abelian group. Fix a point $p \in G$. We define the following:

- For a root class $[\alpha]$, we write $l_{[\alpha]} = \sum_{\xi \in [\alpha]} \xi$ be the sum of all roots in the equivalence class. Let R_G be the set of all $l_{[\alpha]}$'s for $[\alpha]$ ranging over all root classes of G , and G_R for the group of linear maps that leaves R_G invariant. As R_G is finite, G_R is a subgroup of $O(n)$.
- For $\alpha \in \Delta$ a root, we write $\alpha_0 \in \mathbf{A}^*$ for the positive multiple of $l_{[\alpha]}$ of unit norm with respect to the usual Euclidean inner product on $\mathbf{A}$ and $\vec{v}_{[\alpha]} \in \mathbf{A}$ for the dual of α_0 .

- Given $\vec{v} \in \mathbf{A}$, we define

$$\begin{aligned} W_{\vec{v}}^+ &= \oplus_{\Xi(\vec{v}) > 0} V_{\Xi} \\ W_{\vec{v}}^- &= \oplus_{\Xi(\vec{v}) < 0} V_{\Xi} \\ W_{\vec{v}}^0 &= \oplus_{\Xi(\vec{v}) = 0} V_{\Xi} \end{aligned}$$

We say a vector $\vec{v} \in \mathbf{A}$ is *regular* if $\Xi(\vec{v}) \neq 0$ for all roots Ξ , and a linear functional $\ell \in \mathbf{A}^*$ is regular if its dual $\vec{v}_{\ell}$ is regular.

- Let $\ell \in \mathbf{A}^*$, we define $W_{\ell}^+, W_{\ell}^-, W_{\ell}^0$, as $W_{\vec{v}_{\ell}}^+, W_{\vec{v}_{\ell}}^-, W_{\vec{v}_{\ell}}^0$ respectively, where $\vec{v}_{\ell} \in \mathbf{A}$ is the dual of ℓ .
- By *the walls based at p* , we mean the set $p \bigcup_{\Xi} \ker(\Xi)$.
- The root kernels partition the unit sphere in $\mathbf{A}$ into convex subsets called *chambers*. For vectors $\vec{u}, \vec{v}$ in the interior of the same chamber $\mathfrak{b}$, $W_{\vec{u}}^+ = W_{\vec{v}}^+$, and we define $W_{\mathfrak{b}}^+$ for this common subspace of $\mathbf{H}$ and $W_{\mathfrak{b}}^-$ for its complement in $\mathbf{H}$ according to the root space decomposition, so that $\mathbf{H} = W_{\mathfrak{b}}^+ \oplus W_{\mathfrak{b}}^-$.
- By a *geodesic segment through p* , we mean a set $p\overline{AB}$, where $\overline{AB}$ is a directed line segment in $\mathbf{A}$. By direction of a directed line segment in Euclidean space, we mean a unit vector with respect to the usual Euclidean metric, and by direction of $p\overline{AB}$ we mean the direction of $\overline{AB}$.
- For $i = 2, 3, \dots, \text{rank}(G) - 1$, by *i -hyperplane through p* , we mean a set pS , where $S \subset \mathbf{A}$ is an i -dimensional linear subspace or an intersection between an i -dimensional linear subspace with a convex set.
- Let $\pi_A : G \longrightarrow \mathbf{A}$ be the projection onto the $\mathbf{A}$ factor as $(\mathbf{x}, \mathbf{t}) \mapsto \mathbf{t}$.
- For each root α_i , define $\pi_{\alpha_i} : G \longrightarrow V_{\alpha_i} \rtimes \langle \vec{v}_{\alpha_i} \rangle$ as $(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_{|\Delta|})\mathbf{t} \mapsto (\mathbf{x}_i, \alpha_i(\mathbf{t})\vec{v}_{\alpha_i})$. We refer to negatively curved spaces $V_{\alpha_i} \rtimes \langle \vec{v}_{\alpha_i} \rangle$ or $V_{[\alpha]} \rtimes \langle \vec{v}_{\alpha} \rangle$ as weight (or root) hyperbolic spaces.
- For a regular vector $\vec{v} \in \mathbf{A}$, we define $\Pi_{\vec{v}} : G \longrightarrow \mathbf{H} \rtimes \langle \vec{v} \rangle$ as $(\mathbf{x}, \mathbf{t}) \mapsto (\mathbf{x}, \langle \vec{v}, \mathbf{t} \rangle \vec{v})$, where $\langle \cdot, \cdot \rangle$ is the standard inner product on $\mathbf{A}$. In the rank 1 space $\mathbf{H} \rtimes \langle \vec{v} \rangle$, the height function is given as $\mathbf{H} \rtimes \langle \vec{v} \rangle \xrightarrow{\pi_{\vec{v}}} \langle \vec{v} \rangle$
- For a regular liner functional $\ell \in \mathbf{A}^*$ with unit norm, we define $\Pi_{\ell} : G \rightarrow W_{\ell}^- \rtimes \mathbb{R}\vec{v}_{\ell}$ as $(\mathbf{x}, \mathbf{t}) \mapsto ([\mathbf{x}]_{W_{\ell}^-}, \ell(\mathbf{t})\vec{v}_{\ell})$.

Box associated to a compact convex set Fix a net $\mathbf{n}$ of G . For $\alpha \in \Delta$, let $b(r) \subset V_{\alpha}$ be maximal product of intervals of size r , $[0, r]^{\dim(V_{\alpha})}$. Let $\Omega \subset \mathbf{A}$ be a convex compact set with non-empty interior whose barycenter is the identity of $\mathbf{A}$. We define the *box associated to Ω* , $\mathbf{B}(\Omega)$, as the set $\prod_{j=1}^{\sharp} b(e^{\max(\alpha_j(\Omega))})\Omega$. We write

- $\mathcal{L}(\Omega)[m]$ (or $\mathcal{L}(\mathbf{B}(\Omega))[m]$) for the set of geodesics in $\mathbf{B}(\Omega)$ whose π_A images begin and end at points of $\partial\Omega$ such that the ratio between its length and the diameter of Ω lies in the interval $[1/m, m]$. This is a set of $\mathbf{n}$ approximations of such geodesic segments, so is finite.

- For $i = 2, 3, \dots, n$, write $\mathcal{L}_i(\Omega)[m_i]$ (or $\mathcal{L}_i(\mathbf{B}(\Omega))[m_i]$) for the set of i dimensional hyperplanes in $\mathbf{B}(\Omega)$ such that the ratio between its diameter and the diameter of Ω lies in the interval $[1/m_i, m_i]$. This is a set of $\mathbf{n}$ approximations of such bounded subsets in i -hyperplanes, so again is finite.
- $\mathcal{P}(\Omega)$ (or $\mathcal{P}(\mathbf{B}(\Omega))$) for the set of points in $\mathbf{B}(\Omega)$. That is $\mathcal{P}(\Omega) = \mathbf{B}(\Omega) \cap \mathbf{n}$, so a finite set too.
- Let S be an element of $\bigcup_{i=2}^n \mathcal{L}_i(\Omega) \cup \mathcal{L}(\Omega) \cup \mathcal{P}(\Omega)$. We write $L(S)$, $L_i(S)$ for subset of $\mathcal{L}(\Omega)$, $\mathcal{L}_i(\Omega)$ contained or containing S , and $P(S)$ for the subset of $\mathcal{P}(\Omega)$ contained in S .

The following lemma shows that G is amenable.

Lemma 2.2.1. *Let $\Omega \subset \mathbf{A}$ be compact convex with non-empty interior. Then, $\mathbf{B}(r\Omega)$, $r \rightarrow \infty$ is a Følner sequence. The volume ratio between $N_\epsilon(\partial(\mathbf{B}(r\Omega)))$ and $\mathbf{B}(r\Omega)$ is $O(\epsilon/\text{diam}(\mathbf{B}(r\Omega)))^\dagger$.*

Proof. See Lemma ?? in [P] □

Remark 2.2.1. *The same calculation as above shows that for any set $\tilde{B}$ of the form $\Lambda \rtimes \Omega$, where $\Lambda \subset \mathbf{H}$, $\Omega \subset \mathbf{A}$, the ratio of volumes of $N_\epsilon(\partial\tilde{B})$ and that of $\tilde{B}$ is $O(\epsilon/\text{diam}(\tilde{B}))$.*

3 Shadows, slabs and coarsening

We recall the following from [P].

Theorem 3.1. *Let G, G' be non-degenerate, unimodular, split abelian-by-abelian Lie groups, and $\phi : G \rightarrow G'$ be a (κ, C) quasi-isometry. Given $0 < \delta, \eta < \tilde{\eta} < 1$, there exist numbers $L_0, m > 1, \varrho, \hat{\eta} < 1$ depending on $\delta, \eta, \tilde{\eta}$ and κ, C with the following properties:*

If $\Omega \subset \mathbf{A}$ is a product of intervals of equal size at least mL_0 , then a tiling of $\mathbf{B}(\Omega)$ by isometric copies of $\mathbf{B}(\varrho\Omega)$

$$\mathbf{B}(\Omega) = \bigsqcup_{i \in \mathbf{I}} \mathbf{B}(\omega_i) \sqcup \Upsilon$$

contains a subset $\mathbf{I}_0$ of $\mathbf{I}$ with relative measure at least $1 - \nu$ such that

- (i) *For every $i \in \mathbf{I}_0$, there is a subset $\mathcal{P}^0(\omega_i) \subset \mathcal{P}(\omega_i)$ of relative measure at least $1 - \theta$.*
- (ii) *The restriction $\phi|_{\mathcal{P}^0(\omega_i)}$ is within $\hat{\eta}\text{diam}(\mathbf{B}(\omega_i))$ Hausdorff neighborhood of a standard map $g_i \times f_i$.*

Here, ν, θ and $\hat{\eta}$ all approach zero as $\tilde{\eta}, \delta$ go to zero.

In this section, we focus on a particular standard map $g_i \times f_i$ supported on the subset U_i of a good box $\mathbf{B}(\omega_i)$, $i \in \mathbf{I}_0$. We will first show that the f_i is affine for all ranks. Then we will interpret its the constant and linear parts: the linear part has to come from a finite set related to the geometry of G' , and the constant part depends on measure of certain subsets in $\mathbf{B}(\omega_i)$. We will drop the subscript i from now on.

[†]because the ratio of volumes of $\partial\Omega$ to Ω is roughly $\frac{1}{\text{diam}(\Omega)}$, and $\text{rdiam}(\Omega) = \text{diam}(r\Omega)$

3.1 Definitions

In this subsection we define a list of objects that will be used for the remaining of this section.

Root class half planes A set of the form $p \alpha_0^{-1}[-\infty, c]$ (resp. $p \alpha_0^{-1}[c, \infty]$), where $p \in \mathbf{H}$, $c \in \mathbb{R}$, is called a $[\alpha]$ negative (resp. positive) half plane. We write $\mathcal{H}_{[\alpha]}^-$ (resp. $\mathcal{H}_{[\alpha]}^+$) for the set of $[\alpha]$ negative (resp. positive) half planes. When we refer to a $[\alpha]$ half plane in a bounded set, we mean $p \alpha_0^{-1}([c, d])$, for some $p \in \mathbf{H}$, $c, d \in \mathbb{R}$. We will also say that the length of this $[\alpha]$ half plane is $|c - d|$. (remember here that the domain of roots are is $\mathbb{A}$, not the entire group G)

Upper root boundary We define the *upper boundary of root class* $[\alpha]$, $\partial_{[\alpha]}^+$, as the quotient of $\mathcal{H}_{[\alpha]}^+$ under the equivalence relation of bounded Hausdorff distance.

If two positive $[\alpha]$ half-planes E_p, E_q where $p, q \in \mathbf{H}$, are bounded Hausdorff distance apart, then p, q can only differ by $V_{[\alpha]}$ coordinates. This means each equivalence class can be identified with $V_{[\alpha]}$, and the collection of all equivalence classes, $\partial_{[\alpha]}^+$, can be identified with $\oplus_{[\beta] \neq [\alpha]} V_{[\beta]}$.

Lower root boundary We say two $[\alpha]$ negative half planes H_p, H_q are equivalent if there is a sequence $H_i \in \mathcal{H}_{[\alpha]}^-$ such that $H_0 = H_p, H_q = H_n$ and any two successive H_i 's intersect at an unbounded convex set. This is an equivalence relation because if H_p is equivalent to H_q , and H_q is equivalent to H_r , then concatenation of the sequences used to connect the two pairs is a sequence that connects H_p and H_r . We define the *lower boundary of* $[\alpha]$ $\partial_{[\alpha]}^-$ as the quotient of $\mathcal{H}_{[\alpha]}^-$ under this equivalence relation.

We see that if $H_p, H_q \in \mathcal{H}_{[\alpha]}^-$ based at $p, q \in \mathbf{H}$ have non-empty intersection, then p and q cannot differ by $V_{[\alpha]}$ coordinate. On the other hand, if p and q differ only in some $V_{[\beta]}$ coordinate, where $[\beta] \neq [\alpha]$, then $H_p \cap H_q \neq \emptyset$, so H_p is equivalent to H_q in this case. This way, we see that the equivalence class containing $H_p, p \in \mathbf{H}$ are all those $H_q \in \mathcal{H}_{[\alpha]}^-$, where $q \in \mathbf{H}$ differ from p by some elements of $\oplus_{[\beta] \neq [\alpha]} V_{[\beta]}$, and consequently, $\partial_{[\alpha]}^-$ can be identified with $V_{[\alpha]}$.

Measures on lower root boundaries If $p \in G$, we write $\pi_{[\alpha]}^-(p) \subset \partial_{[\alpha]}^-$ (resp. $\pi_{[\alpha]}^+(p) \subset \partial_{[\alpha]}^+$) for the set of equivalence classes, each containing a minimal negative (resp. positive) $[\alpha]$ half planes through a point that is at most distance ρ away from p , where ρ is the scale of discretization. Since $V_{[\alpha]}$ is the direct sums of V_{Ξ} , where $\Xi \in [\alpha]$, we will write $\pi_{\sigma}^-(p)$, where $\sigma \in [\alpha]$, for the V_{σ} coordinate of $\pi_{[\alpha]}^-(p)$. For $A \subset G$, we write $\pi_{[\alpha]}^*(A) = \bigcup_{p \in A} \pi_{[\alpha]}^*(p)$, where $*$ is $\{+, -\}$.

Since $\partial_{[\alpha]}^-$ is a homogeneous space (the subgroup $V_{[\alpha]} \subset \mathbf{H}$ acts faithfully and transitively on it), it admits a Haar measure. We normalize this measure $|\cdot|$ by requiring that for each $\sigma \in [\alpha]$,

$$|\pi_{\sigma}^-(p)| e^{-\sigma(p)} = 1, \forall p \in G \quad (3)$$

Upper and lower boundaries of a linear functional We call the intersection of half planes corresponding to two perpendicular linear functionals, a quarter plane.

We now define an equivalence relation on $\mathcal{H}_{\ell}^+$ (resp. $\mathcal{H}_{\ell}^-$) as follows. Two positive (resp. negative) ℓ half planes H_p, H_q are equivalent if there is a sequence of $H_i \in \mathcal{H}_{\ell}^+$ such that $H_0 = H_p, H_n = H_q$ and the intersection between any two successive H_i 's does not contain a quarter plane. This is an equivalence relation because if H_p is equivalent to H_q , and H_q is equivalent to H_r , then the concatenation of the sequences used to connect the two pairs is a sequence that establishes

equivalence between H_p and H_r . We see that if the positive (resp. negative) ℓ half planes H_p, H_q based at $p, q \in \mathbf{H}$ are equivalent, then p, q differ by an element of W_ℓ^- . (resp. W_ℓ^+)

We define the *upper boundary* of ℓ , ∂_ℓ^+ , (resp. *lower boundary* of ℓ , ∂_ℓ^-) as the quotient of $\mathcal{H}_\ell^+$ (resp. $\mathcal{H}_\ell^-$) under this equivalence relation. In light of the forgoing discussion, we see that ∂_ℓ^+ (resp. ∂_ℓ^-) can be identified with W_ℓ^+ . (resp. W_ℓ^-)

Measure on upper and lower boundaries of a linear functional Let ℓ be a generic linear functional. Since ∂_ℓ^+ can be identified with W_ℓ^+ which itself is a direct sum of $V_{[\Xi]}$ where $\ell(\vec{v}_\Xi) > 0$, and each of $V_{[\Xi]}$ can be identified with $\partial_{[\Xi]}$, we can identify ∂_ℓ^+ with $\prod_{[\Xi]: \ell(\vec{v}_\Xi) > 0} \partial_{[\Xi]}$, and equip it with the product measures on the constituent root boundaries. The same procedure can be applied to ∂_ℓ^- to turn into a measure space.

If $p \in G$, we write $\pi_\ell^-(p) \subset \partial_\ell^-$ (resp. $\pi_\ell^+(p) \subset \partial_\ell^+$) for the set of equivalence classes, each containing a smallest negative (resp. positive) ℓ half planes through a point that is at most distance ρ away from p , where ρ is the scale of discretization. For $A \subset G$, and $* \in \{+, -\}$, $\pi_\ell^*(A)$ is the union of $\pi_\ell^+(p)$, where p ranges over all points of A .

Branching constant The branching constant $\mathbf{b}_{[\alpha]}$ of root class $[\alpha]$ is the number such that $e^{\mathbf{b}_{[\alpha]}L}$ represents the number of $[\alpha]$ half planes of length L leaving a point. It equals $l_{[\alpha]}/\alpha_0$.

The branching constant $\mathbf{b}_\ell$, of a generic linear functional ℓ , is a number such that $e^{\mathbf{b}_\ell L}$ represents the number of ℓ half planes of length L leaving a point. Its value is given by

$$\mathbf{b}_\ell = \sum_{\sigma: \sigma(\vec{v}_\ell) > 0} \sigma(\vec{v}_\ell) = \sum_{\sigma: \sigma(\vec{v}_\ell) < 0} \sigma(\vec{v}_\ell) \quad (4)$$

If $\tilde{\ell}$ is a linear functional whose norm is not 1, we will write $\mathbf{b}_{\tilde{\ell}}$ for $\mathbf{b}_{\tilde{\ell}/\|\tilde{\ell}\|}$.

Distances on lower root boundaries Given $p, q \in \partial_{[\alpha]}^- \sim V_{[\alpha]}$, let $t_{p,q}$ be the minimal $t \in \mathbb{R}$ such that there exists negative $[\alpha]$ half planes in the equivalence class of p and q that are distance 1 (or ρ if the scale of discretization is not 1) at sets whose π_A projection is $\ell_{[\alpha]}^{-1}(t)$.

Fix a positive number c , we define a *pseudo distance* $D_{[\alpha]}$ between p, q as

$$D_{[\alpha]}(p, q) = e^{ct_{p,q}}$$

Different choice of c leads to quasi-symmetric equivalent metric. In this way, the space $(\partial_{[\alpha]}^-, D_{[\alpha]})$ becomes those whose quasi-conformal maps are studied in [D].

Shadows, slabs Using the same root class as before, we define any subset H of a left translate of $\oplus_{[\beta] \neq [\alpha]} V_{[\beta]} \rtimes \ker(\alpha_0)$ a $[\alpha]$ *block*. Note that for any element $\sigma \in [\alpha]$, $\sigma(H)$ is well-defined. For $\rho > 1$, we define the ρ -*shadow* of H , $Sh(H, \rho)$, as the union of smallest negative $[\alpha]$ half planes containing a point in $N_\rho H$. For $h_2 < h_1 < \alpha_0(H)$, we define a *slab* of H , denoted by $Sl_2^1(H)$ as the intersection between $Sh(H, \rho)$ with $\alpha_0^{-1}([h_2, h_1])$. That is, it's the subset of $Sh(H, \rho)$ that whose α_0 values lies in between h_2 and h_1 .

Generalized slabs For $E_- \subset \partial_{[\alpha]}^-$, $E^+ \subset \partial_{[\alpha]}^+$, $K \subset \ker(\alpha_0)$, $h_2 < h_1$, we call a set

$$S(E_-, E^+, K, h_2, h_1) = \{(\mathbf{x}_{[\alpha]}, (\mathbf{x}_{[\beta]})_{[\beta] \neq [\alpha]}, \mathbf{t}) : \mathbf{t} \in [h_2, h_1]K, \mathbf{x}_{[\alpha]} \in E_-, (\mathbf{x}_{[\beta]})_{[\beta] \neq [\alpha]} \in E^+\}$$

a *generalized $[\alpha]$ slab*. This generalizes the definition of slabs defined in the previous paragraph.

Coarsening We define a process *coarsening* as follows. For $h \in \mathbb{R}$, $E^+ \subset \partial_{[\alpha]}^+$, the *coarsening of E^+ by h* , $\mathcal{C}_h(E^+)$ is defined as the subset of $\partial_{[\alpha]}^+$ consisting of those equivalence classes that contains a positive $[\alpha]$ half plane that has a non-empty intersection with a positive $[\alpha]$ half plane whose equivalence class belongs to E^+ at a set whose π_A projection belongs to $\alpha_0^{-1}[h, \infty]$.

Similarly, for $E_- \subset \partial_{[\alpha]}^-$ a subset of the lower boundary, the coarsening of E_- by h , $\mathcal{C}_h^{E_-}$, is defined to be those equivalence classes of negative $[\alpha]$ half planes that contains an element which intersects non-empty with a negative $[\alpha]$ half plane whose equivalence class belongs to E_- at a set whose π_A projection is a subset of $\alpha_0^{-1}([-\infty, h])^\dagger$.

Observe that as long as $h_3 \leq h_2$, $h_4 \geq h_1$, we have

$$S(E_-, E^+, h_2, h_1) = S(\mathcal{C}_{h_3}(E_-), \mathcal{C}_{h_4}(E^+), h_2, h_1)$$

Lemma 3.1.1. *The number of $[\alpha]$ planes in $S = S(\mathcal{C}_{h_3}(E_-), \mathcal{C}_{h_4}(E^+), K, h_2, h_1)$ is comparable to*

$$\frac{\text{Vol}(S)}{|K|(h_1 - h_2)} e^{\mathbf{b}_{[\alpha]}(h_1 - h_2)}$$

That is, it is compatible to the area of the cross-section times $e^{\mathbf{b}_{[\alpha]}(h_1 - h_2)}$.

Proof. Counting the number of $[\alpha]$ half planes really means you count the number of geodesics in $V_{[\alpha]} \times \mathbb{R}(\vec{v}_{[\alpha]})$. In this way, we see that the first term of the product is the size of a cross section in this projection image, and the product is the no. of geodesics in the range. $\square$

3.2 Improving almost product map

Recall our setting from Theorem 3.1.

$$\phi : \mathbf{B}(\omega) \rightarrow G'$$

is a quasi-isometry such that on a subset $U_* \subset \mathbf{B}(\omega)$ of relative measure at least $1 - \theta$, the restriction $\phi|_{U_*}$, is within $\hat{\eta} \text{diam}(\mathbf{B}(\omega))$ of a standard map $\hat{\phi} = g \times f$.

If $U_* = \mathbf{B}(\omega)$, then it is clear that the image of a slab is a slab, and image of a generalized slab is also a generalized slab. However as U_* is generally not $\mathbf{B}(\omega)$, it is not clear that for a $[\alpha]$ block H , and $h_2 < h_1 < \alpha_0(H)$, there is an obvious relation between $\phi(Sl_2^1(H))$ and $\hat{Sl}_2^1(H)$, which is defined as $S(g(\pi_{[\alpha]}^-(H)), g(\pi_{[\alpha]}^+(H)), f(\pi_A(H) \times [h_2, h_1]))$, other than

$$\phi(Sl_2^1(H) \cap U_*) \subset N_{\hat{\eta} \text{diam}(\mathbf{B}(\omega))} \hat{\phi}(Sl_2^1(H) \cap U_*) \subset N_{\hat{\eta} \text{diam}(\mathbf{B}(\omega))} \hat{Sl}_2^1(H)$$

We will show that by restricting to certain subset of $Sl_2^1(H)$ whose coarsened version lies almost entirely in U_* , a reversal inclusion can be obtained.

To this end, fix a $[\alpha]$ block H in $\mathbf{B}(\omega)$, and let $h = \alpha_0(H)$. As f preserves foliations by root kernels, the f image of $\ker(\alpha_0)$ is the kernel of $\Xi_0 \in \mathbf{A}^*$ for some root class $[\Xi]$. We identify $\mathbf{A}/\ker(\alpha_0)$ with $\mathbb{R}$ by taking the α_0 value of a coset, and do so similarly for $\mathbf{A}/\ker(\Xi_0)$. Then f

[†]This is the same as the set of points in $\partial_{[\alpha]}^-$ that is distance e^h from a point in E_- .

induces a map $q : \mathbb{R} \rightarrow \mathbb{R}$ by sending t to the Ξ_0 value of $f(t \ker(\alpha_0))$. When the rank of G is 1, this is just f .

Now, fix heights $h_2 < h_1 < h$, and define

$$\hat{Sl}_2^1(H) = S(\mathcal{C}_{q(h_1)}g(\pi_{[\alpha]}^-(H)), \mathcal{C}_{q(h_2)}g(\pi_{[\alpha]}^+(H)), f(\pi_A(H) \times [h_2, h_1]))$$

In the rest of this section, we write B for the box $\mathbf{B}(\omega) = \left(\prod_{[\sigma]} A_{[\sigma]}\right) \omega$ where $A_{[\sigma]} \subset V_{[\sigma]}$, $\omega \subset \mathbf{A}$.

We refer to the parameters $\delta, \eta, \tilde{\eta}$ in Theorem 3.1 as ‘initial data’, and show now that most slabs inherit the property that ‘intersection with U_* possess relative large measure’ For the remaining subsections, whenever $\gg$ appears, it means the ratio of the two quantities is more than 2κ .

Lemma 3.2.1. *Given $0 < \beta \ll \beta' \ll 1$, there exist constants $c_1, c_2 \ll 1$ depending on our initial data and a subset $E_{**} \subset \partial_{[\alpha]}^-(B)$ of relative large measure such that whenever H is a $[\alpha]$ block which is at least $2\kappa\beta' \text{diam}(\mathbf{B}(\omega))$ away from ∂B and $\pi_{[\alpha]}^-(H) \cap E_{**} \neq \emptyset$, then $|h_1(H) - h_2(H)| |\pi_A(H)| > \beta |\omega|$ implies*

$$|Sl_2^1(H) \cap U_*| \geq (1 - c_2) |Sl_2^1(H)| \quad (5)$$

Proof. Let c_2 be a constant to be chosen later. Let $E_1 \subset \partial_{[\alpha]}^-(B)$ be the subset such that for $x \in E_1$, there exists a $[\alpha]$ block H_x such that $x \in I_x = \pi_{[\alpha]}^-(H_x)$ and equation (5) fails. Then we have a cover of E_1 by intervals I_x . By Vitali covering there is a subset of I_k ’s such that $\sum_k |I_k| \geq 1/5 |E_1|$, and whose elements are strongly disjoint i.e. for $j \neq k$, $d(I_j, I_k) \geq 1/2 \max(|I_j|, |I_k|)$, which means that the corresponding $[\alpha]$ block H_k ’s are also disjoint. By construction $|Sl_2^1(H_k) \cap U_*^c| \geq c_2 |Sl_2^1(H_k)|$.

Summing over k yields

$$|B \cap U_*^c| \geq c_2 \sum_k |Sl_2^1(H_k)| \geq \frac{c_2}{2} \sum_k |h_1(H_k) - h_2(H_k)| |\pi_A(H_k)| |I_k| |\pi_{[\alpha]}^+(H_k)|$$

As $|\pi_{[\alpha]}^+(H_k)| = |\pi_{[\alpha]}^+(B)| = \prod_{[\sigma] \neq [\alpha]} |A_{[\sigma]}|$, and $|B \cap U_*^c| \leq \theta |\omega| |\pi_{[\alpha]}^-(B)| |\pi_{[\alpha]}^+(B)|$, we obtain

$$|E_1| \leq 5 \sum_k |I_k| \leq \frac{10\theta}{\beta c_2} |\partial_{[\alpha]}^-(B)|$$

Now choose c_2 appropriately. □

Note that as $\hat{\phi}$ is only partially defined, the definition of $\hat{Sl}_2^1(H)$ sort of ‘fills up’ $\hat{\phi}(Sl_2^1(H))$. Therefore it is not clear that most of $\hat{Sl}_2^1(H)$ lie close to the $\phi(U_*)$. We now introduce some notations needed to describe a subset of $\hat{Sl}_2^1(H)$ which can be controlled more easily. Given a set $D \subset B$ and $A_{[\Xi]} \subset \pi_{[\Xi]}^+(B)$, we write $A_{[\Xi]} \cap D$ for the intersection between $A_{[\Xi]}$ and $\pi_{[\Xi]}^+(D)$. We then define

$$\tilde{Sl}_2^1(H, D) := S(\mathcal{C}_{q(h_1)}(g(\pi_{[\alpha]}^-(H))), \mathcal{C}_{q(h_2)}(g(\pi_{[\alpha]}^+(H) \cap D)), f(\pi_A(H) \times [h_2, h_1]))$$

This next lemma says that for certain ‘slimmed down’ version of $\hat{Sl}_2^1(H)$, $\tilde{Sl}_2^1(H, D)$, most of them lies in $\phi(U_*)$.

Lemma 3.2.2. *Given $\hat{\eta} < \beta \ll \beta' \ll \beta'' < 1$, there exist constants c_3, c_4 depending on our initial data and a subset $E_* \subset \pi_{[\alpha]}^-(B)$ of relative large measure with the following properties.*

Let H_0 be a $[\alpha]$ block in B such that

- a. The distance between H_0 and ∂B is at least $4\kappa\beta'' \text{diam}(\mathbf{B}(\omega))$.*
- b. The intersection $\pi_{[\alpha]}^-(H_0) \cap E_*$ is not empty.*

Suppose H is a $[\alpha]$ block in $S = U_ \cap S(\pi_{[\alpha]}^-(H_0), \pi_{[\alpha]}^+(H_0), \pi_A(H_0), \ell_{[\alpha]}(H_0), \ell_{[\alpha]}(H_0) + \beta'' \text{diam}(\mathbf{B}(\omega)))$ such that $\pi_{[\alpha]}^-(H)$ has non-empty intersection with E_* . Then, $\beta'|\omega| > |h_1(H) - h_2(H)| |\pi_A(H)| > \beta|\omega|$ implies*

$$\left| \tilde{S}l_2^1(H) \cap \phi(U_*) \right| \geq (1 - c_4) \left| \tilde{S}l_2^1(H) \right| \quad (6)$$

where $\tilde{S}l_2^1(H) = \tilde{S}l_2^1(H, S)$, and c_4 approaches zero as our initial data approach zero's.

Proof. The set E_* will be constructed from E_{**} that appeared in Lemma 3.2.1.

We first show that $\tilde{S}l_2^1(H) \subset \phi(B)$. Recall that H_0 is more than $4\kappa^2\beta'' \text{diam}(\mathbf{B}(\omega))$ away from the boundary of B . This means that S is also more than $4\kappa^2\beta'' \text{diam}(\mathbf{B}(\omega))$ away from ∂B . By definition

$$\tilde{S}l_2^1(H) = S(\mathcal{C}_{q(h_1)}(g(\pi_{[\alpha]}^-(H))), \mathcal{C}_{q(h_2)}g(\pi_{[\alpha]}^+(H) \cap S), f(\pi_A(H) \times [h_2(H), h_1(H)]))$$

By assumption, $|h_1(H) - h_2(H)| \leq \beta' \frac{|\omega|}{|\pi_A(H)|}$, so $|h_1 - h_2| \leq \beta' \text{diam}(\mathbf{B}(\omega))$. Take $q_0 \in \tilde{S}l_2^1(H)$. Then q_0 is no further than $\beta' \text{diam}(\mathbf{B}(\omega))$ away from a point q in the generalized $[\alpha]$ slab $S(g(\pi_{[\alpha]}^-(H)), g(\pi_{[\alpha]}^+(H) \cap S), f(\pi_A(H) \times [h_2(H), h_1(H)]))$. This means that there is $p \in S \subset U_*$ such that $\pi_{[\alpha]}^+(q) = \pi_{[\alpha]}^+(\hat{\phi}(p))$, so $d(q, \hat{\phi}(p)) \leq \beta' \text{diam}(\mathbf{B}(\omega))$.

By definition, $\hat{\phi}(p)$ lies on a $[\Xi]$ half plane that is $\hat{\eta} \text{diam}(\mathbf{B}(\omega))$ away from the image of a $[\alpha]$ half-plane containing p . This means $d(q, \phi(S)) \leq 2\beta' \text{diam}(\mathbf{B}(\omega))$, and therefore $d(q_0, \phi(S)) \leq 3\beta' \text{diam}(\mathbf{B}(\omega))$. Since $d(S, \partial B) > 4\beta'' \text{diam}(\mathbf{B}(\omega))$, it follows that $q_0 \in \phi(B)$.

Let c_3 be a constant to be chosen later. Let $E_2 \subset \partial_{[\alpha]}^-(B) - E_1$ be such that for $x \in E_2$ there is a $[\alpha]$ block H_x such that $x \in I_x = \pi_{[\alpha]}^-(H_x)$ and equation (6) fails. Thus we have a cover of E_2 by intervals I_x . By Vitali covering, we can find a subset of I_k 's such that the inequality opposite to above holds for each H_k (instead of H) and $\sum_k |I_k| \geq (1/5)|E_2|$, and that I_k 's are strongly disjoint¹. (That is, for $j \neq k$, $d(I_j, I_k) \geq 1/2 \max(|I_j|, |I_k|)$). In particular, this means $\tilde{S}l_2^1(H_k)$'s are disjoint as well.

We now claim that

$$\phi(Sh(H_k, O(1))^c \cap U_*) \cap \tilde{S}l_2^1(H_k) = \emptyset \quad (7)$$

Suppose not, then there is a $p \in Sh(H_k, O(\rho_1))^c \cap U_*$ such that $\phi(p) \in \tilde{S}l_2^1(H_k)$. The latter means that $\pi_{[\alpha]}^-(\phi(p)) \in \mathcal{C}_{q(h_1(H))}g(\pi_{[\alpha]}^-(H_k))$, so there is a $p' \in Sh(H_k, O(1)) \cap U_*$ such that

$$t_{\pi_{[\Xi]}^-(\phi(p)), \pi_{[\Xi]}^-(\hat{\phi}(p'))} \leq q(h_1(H_k)) + O(\hat{\eta} \text{diam}(\mathbf{B}(\omega))) \quad (8)$$

¹Note that we cannot proceed in the same manner as Lemma 3.2.1 from this point on because the $\partial_{[\alpha]}^+(S)$ is not a full set: it's the intersection of this with something else, so we need to do something else first.

However the fact that $p \in Sh(H_k, O(1))^c$ and $p' \in Sh(H_k, O(1))$ means that for

$$t_{\pi_{[\alpha]}^-(p), \pi_{[\alpha]}^-(p')} > \alpha_0(H_k) + O(1)$$

which in turns means that

$$t_{\pi_{[\Xi]}^-(\phi(p)), \pi_{[\Xi]}^-(\hat{\phi}(p'))} > q(\alpha_0(H_k)) + O(\hat{\eta}diam(\mathbf{B}(\omega))) > q(h_1(H_k)) + O(\hat{\eta}diam(\mathbf{B}(\omega)))$$

contradicting (8). So equation (7) holds.

Next we show that

$$\tilde{Sl}_2^1(H_k) \cap N_{O(\hat{\eta}diam(\mathbf{B}(\omega)))} \hat{\phi}(Sl_2^1(H_k))^c \subset \phi(U_*^c) \quad (9)$$

Since U_*^c has relative small measure in B , this means that the measure of left hand side over all k 's from the strongly disjoint family remains small, making the measure of $\sum_k N_{O(\hat{\eta}diam(\mathbf{B}(\omega)))} \hat{\phi}(Sl_2^1(H_k))$ a lower bound for $\sum_k |\tilde{Sl}_2^1(H_k)|$.

Suppose (9) is not true. Then there is a $p \in \tilde{Sl}_2^1(H_k)$, such that $p \in N_{O(\hat{\eta}diam(\mathbf{B}(\omega)))} \hat{\phi}(Sl_2^1(H_k))^c$ and $p \in \phi(U_*)$. However, by equation (7), the last two conditions means that $p \notin \tilde{Sl}_2^1(H_k)$, which contradicts assumption that $p \in \tilde{Sl}_2^1(H_k)$.

By choice, $\tilde{Sl}_2^1(H_k)$'s satisfy the inequality opposite to equation (6), so $|\tilde{Sl}_2^1(H_k) \cap \phi(U_*^c)| \geq c_4 |\tilde{Sl}_2^1(H_k)|$. As all the $\tilde{Sl}_2^1(H_k)$'s are disjoint, summing over k yields

$$\begin{aligned} |\phi(U_*^c)| &\geq c_4 \sum_k |\tilde{Sl}_2^1(H_k)| \geq c_4(1-\theta) \sum_k N_{O(\hat{\eta}diam(\mathbf{B}(\omega)))} \hat{\phi}(Sl_2^1(H_k)) \\ &\geq c_4(1-\theta)(1-O(\hat{\eta}))(1-c_2) \sum_k |Sl_2^1(H_k)| \\ &\geq c_4(1-\theta)(1-O(\hat{\eta}))(1-c_2) \sum_k |\pi_A(H_k)| |h_1(H_k) - h_2(H_k)| |I_k| |\pi_{[\alpha]}^+(H_k)| \end{aligned}$$

The second line comes from Lemma 3.2.1 and the fact that the volume ratio between ϵ neighborhood of $Sl_2^1(H)$ and $Sl_2^1(H)$ is $1 + O(\epsilon)$ in the second line, as noted in Remark 2.2.1.

Since $\pi_{[\alpha]}^+(H_k) = \pi_{[\alpha]}^+(B)$, and $|\phi(U_*^c)| \leq \theta |\omega| |\pi_{[\alpha]}^+(B)| |\pi_{[\alpha]}^-(B)|$, we obtain

$$|E_2| \leq 5 \sum_k |I_k| \leq \frac{5\theta}{c_4 \beta (1-\theta)(1-O(\hat{\eta}))(1-c_2)} |\pi_{[\alpha]}^-(B)|$$

□

Corollary 3.2.1. *Let H be a $[\alpha]$ block in $\mathbf{B}(\omega)$ satisfying the hypothesis of Lemma 3.2.1 and 3.2.2, and S as in Lemma 3.2.2. Let $w_1, w_2 \in \mathbb{R}$ be such that $\beta'|\omega| \geq |w_1 - w_2| |\pi_A(H)| \geq \beta|\omega|$. Then*

$$|\mathcal{C}_{w_1}(g(\pi_{[\alpha]}^-(H)))| |\mathcal{C}_{w_2}(g(\pi_{[\alpha]}^+(H) \cap S))| \geq d |\pi_{[\alpha]}^-(H)| |\pi_{[\alpha]}^+(H)| \quad (10)$$

and

$$|\mathcal{C}_{w_1}(g(\pi_{[\alpha]}^-(H)))| |\mathcal{C}_{w_2}(g(\pi_{[\alpha]}^+(H) \cap S))| \leq b |\pi_{[\alpha]}^-(H)| |\pi_{[\alpha]}^+(H)| \quad (11)$$

where d and b depend only on κ, C .

Proof. Note that from the structure of U and the fact that ϕ is a quasi-isometry, it follows that for $z_1, z_2 \in \pi_A(\mathbf{B}(\omega))$ we have

$$\frac{1}{2\kappa} |z_1 - z_2| - \hat{\eta} \text{diam}(\mathbf{B}(\omega)) \leq |q(z_1) - q(z_2)| \leq 2\kappa |z_1 - z_2| + \hat{\eta} \text{diam}(\mathbf{B}(\Omega))$$

This means that q is essentially monotone, so there exists h_1, h_2 such that $q(h_1(H)) = w_1$, and $q(h_2(H)) = w_2$. We now apply Lemma 3.2.1 and 3.2.2 to the resulting $Sl_2^1(H)$ and $\tilde{Sl}_2^1(H)$.

By equation (6) in Lemma 3.2.2, we know that $|\tilde{Sl}_2^1(H) \cap \phi(U_*)| \geq (1 - c_4) |\tilde{Sl}_2^1(H)|$. The structure of a standard map means that the ratio of measures of $\tilde{Sl}_2^1(H) \cap \phi(U_*)$ to that of $Sl_2^1(H) \cap U_*$ lies in $[1/2\kappa, 2\kappa]$. These facts together shows that

$$(1 - c_4) |\tilde{Sl}_2^1(H)| \leq |\tilde{Sl}_2^1(H) \cap \phi(U_*)| \leq 2\kappa |Sl_2^1(H) \cap U_*| \leq 2\kappa |Sl_2^1(H)|$$

Similarly,

$$(1 - c_2) |Sl_2^1(H)| \leq |Sl_2^1(H) \cap U_*| \leq 2\kappa |\tilde{Sl}_2^1(H) \cap \phi(U_*)| \leq 2\kappa |\tilde{Sl}_2^1(H)|$$

The claims now follow from the following volume formula.

$$|Sl_2^1(H)| = |\pi_{[\alpha]}^-(H)| |\pi_{[\alpha]}^+(H)| |\pi_A(H)| |h_2 - h_1|$$

$$|\tilde{Sl}_2^1(H)| = |\mathcal{C}_{q(h_1)} g(\pi_{[\alpha]}^-(H))| |\mathcal{C}_{q(h_2)} g(\pi_{[\alpha]}^+(H))| |f(\pi_A(H))| |q(h_1) - q(h_2)|$$

□

3.3 The constant part of f

Continuing with the same notations from the previous subsection, we show in this subsection, that $q : \mathbf{A}/\ker(\ell_{[\alpha]}) \rightarrow \mathbf{A}/\ker(\ell_{[\Xi]})$ is affine and compute its constant term. To compute the the constant term of this affine map, we make use of Corollary 3.2.1, and the property that the standard map $\hat{\phi}$ roughly preserves the number of root class half planes.

Lemma 3.3.1. *Let H be a $[\alpha]$ block in B . Let $\mathcal{F}, \tilde{\mathcal{F}}$ denote the set of (maximal) $[\alpha], [\Xi]$ half planes in $Sl_2^1(H)$ and $\tilde{Sl}_2^1(H)$ respectively. Then*

$$\log |\mathcal{F}| = \log |\tilde{\mathcal{F}}| + O(\hat{\eta} \text{diam}(B))$$

Proof. The claim is base on the fact that there is a bijection between $Sl_2^1(H) \cap U_*$ and $\tilde{Sl}_2^1(H) \cap \phi(U_*)$.

Explicitly, let $\mathcal{F}'$ be the set of $[\alpha]$ half planes in $Sl_2^1(H)$ that are more than $O(\hat{\eta} \text{diam}(B))$ away from ∂B , and spend at least $1 - \sqrt{c_2}$ fraction of their measure in U_* . Then $\mathcal{F}'$ has a relative large measure in $\mathcal{F}$.

For each $\gamma \in \mathcal{F}'$ there is a $[\Xi]$ half plane $\hat{\gamma} \in \tilde{\mathcal{F}}$ such that $\phi(\gamma \cap U_*)$ is within $\hat{\eta} \text{diam}(B)$ of $\hat{\gamma}$. We define $\psi(\gamma) = \hat{\gamma}$. Note ψ is at most $e^{\hat{\eta} \text{diam}(B) + \sqrt{c_2} \text{diam}(B)}$ to one since two $[\alpha]$ planes with the same $\hat{\phi}$ image must be within $\hat{\eta} \text{diam}(B)$ of each other whenever they are in U_* . Inverse of ψ is defined similarly. □

Lemma 3.3.2. *For all $z_1, z_2 \in [z_{bot}, z_{top}]$, where $z_{top} = \max \alpha_0(B)$, and $z_{bot} = \min \alpha_0(B)$.*

$$q(z_1) - q(z_2) = \frac{\mathbf{b}_{[\alpha]}}{\mathbf{b}_{[\Xi]}}(z_1 - z_2) + O(\hat{\eta} \text{diam}(B))$$

where $[\Xi] = f_*[\alpha]$

Proof. It is sufficient to check this for a $[\alpha]$ block H , and h_1, h_2 satisfying the hypothesis of Lemma 3.2.1 and 3.2.2. Let $\mathcal{F}, \tilde{\mathcal{F}}$ denote the set of $[\alpha], [\Xi]$ half planes in $Sl_2^1(H)$ and $\tilde{Sl}_2^1(H)$ respectively. The number of $[\alpha]$ half planes in $Sl_2^1(H)$ is

$$|\pi_{[\alpha]}^-(H)| |\pi_{[\alpha]}^+(H)| e^{\mathbf{b}_{[\alpha]}(h_1 - h_2)}$$

while the number of $[\Xi]$ half planes is

$$|\mathcal{C}_{q(h_1)}g(\pi_{[\alpha]}^-(H))| |\mathcal{C}_{q(h_2)}g(\pi_{[\alpha]}^+(H) \cap S)| e^{\mathbf{b}_{[\Xi]}(q(h_1) - q(h_2))}$$

By Lemma 3.3.1,

$$|\mathcal{C}_{q(h_1)}g(\pi_{[\alpha]}^-(H))| |\mathcal{C}_{q(h_2)}g(\pi_{[\alpha]}^+(H) \cap S)| e^{\mathbf{b}_{[\Xi]}(q(h_1) - q(h_2))} = |\pi_{[\alpha]}^-(H)| |\pi_{[\alpha]}^+(H)| e^{\mathbf{b}_{[\alpha]}(h_1 - h_2)} e^{\hat{\eta} \text{diam}(B)}$$

After simplifying,

$$q(h_1) - q(h_2) = \frac{\mathbf{b}_{[\alpha]}}{\mathbf{b}_{[\Xi]}}(h_1 - h_2) + \frac{1}{\mathbf{b}_{[\Xi]}} \log \frac{|\pi_{[\alpha]}^-(H)| |\pi_{[\alpha]}^+(H)|}{|\mathcal{C}_{q(h_1)}g(\pi_{[\alpha]}^-(H))| |\mathcal{C}_{q(h_2)}g(\pi_{[\alpha]}^+(H) \cap S)|} + \hat{\eta} \text{diam}(B)$$

The claim now follows because by Corollary 3.2.1,

$$\frac{|\pi_{[\alpha]}^-(H)| |\pi_{[\alpha]}^+(H)|}{|\mathcal{C}_{q(h_1)}g(\pi_{[\alpha]}^-(H))| |\mathcal{C}_{q(h_2)}g(\pi_{[\alpha]}^+(H) \cap S)|} = O(1)$$

□

Lemma 3.3.3. *The $\mathbf{A}'$ part of a standard map is affine. Its natural action sends R_G to $R_{G'}$, hence can only take on one of finitely many possibilities.*

Proof. When G has exact $\text{rank}(G)$ many root kernels, every point of $\mathbf{A}$ and $\mathbf{A}'$ is uniquely determined by the intersection of $\text{rank}(G)$ many translates of root kernels, so Lemma 3.3.2 shows that f is affine and takes foliations by root kernels of G to that of G' .

Let A denote for the linear part of f , and σ be the permutation that f induces on the root classes. The existence of a standard map $O(\hat{\eta} \text{diam}(B))$ away from a quasi-isometry means that for $\vec{u}$ ranges over a large subset of $\mathbb{S}^{n-1}$, $[\Xi]$ a root class, then $l_{[\Xi]}(\vec{u}) > 0$ if and only if $l_{\sigma([\Xi])}(A(\vec{u})) > 0$, and $l_{[\Xi]}(\vec{u}) = 0$ if and only if $l_{\sigma([\Xi])}(A(\vec{u})) = 0^\dagger$. So $l_{\sigma([\Xi])} \circ A = c_{[\Xi]} l_{[\Xi]}$ for some $c_{[\Xi]} > 0$.

Since f is affine, the push-forward of $\ell \in \mathbf{A}^*$, $f_*(\ell) = \ell \circ A^{-1}$, is an element of $\mathbf{A}'^*$. Take ℓ a regular linear functional of unit norm, and H an $[\alpha]$ block. Then, inside of $Sl_2^1(H)$, the number of maximal sets of the form $p \ell^{-1}[c, d]$, $p \in \mathbf{H}$, $d - c = L$ is

[†]Note that in general, $l_{[\Xi]} \circ A^{-1} = c l_{\sigma([\Xi])}$ for some positive c , not necessarily 1. All that we know is that A induces a permutation on the root classes, not on the set R_G .

$$|\pi_{[\alpha]}^-(H)| |\pi_{[\alpha]}^+(H)| e^{L \sum_{[\Xi]: \Xi(\vec{v}_\ell) > 0} l_{[\Xi]}(\vec{v}_\ell)}$$

while the number of $f_*\ell$ half planes in $\tilde{S}l_2^1(H)$ is

$$|\mathcal{C}_{q(h_1)}g(\pi_{[\alpha]}^-(H))| |\mathcal{C}_{q(h_2)}g(\pi_{[\alpha]}^+(H) \cap S)| e^{L \sum_{[\Xi]: \Xi(\vec{v}_\ell) > 0} l_{\sigma([\Xi])}(A\vec{v}_\ell)}$$

Simplifying using Lemma 3.3.1 and Corollary 3.2.1 yields

$$\sum_{[\Xi]: \Xi(\vec{v}_\ell) > 0} l_{\sigma([\Xi])}(A\vec{v}_\ell) = \sum_{[\Xi]: \Xi(\vec{v}_\ell) > 0} l_{[\Xi]}(\vec{v}_\ell) + \hat{\eta} \text{diam}(B)$$

Since the map from S^{n-1} to disjoint union of root classes defined by sending $\vec{v}$ to $\{[\Xi] : \Xi(\vec{v}) > 0\} \sqcup \{[\beta] : \beta(\vec{v}) < 0\}$ is constant on chambers, and each chamber contains a basis of $\mathbf{A}$, we conclude that up to an error of $\hat{\eta} \text{diam}(\mathbf{B}(\omega))$,

$$\sum_{[\Xi]: \Xi(\vec{v}_\ell) > 0} l_{[\Xi]} = \sum_{[\Xi]: \Xi(\vec{v}_\ell) > 0} l_{\sigma([\Xi])} \circ A = \sum_{[\Xi]: \Xi(\vec{v}_\ell) > 0} c_{[\Xi]} l_{[\Xi]}$$

In other words, we have two equations:

$$\begin{aligned} \sum_{[\alpha]: \alpha(\vec{v}) > 0} (1 - c_{[\alpha]}) l_{[\alpha]} &= 0 \\ \sum_{[\beta]: \beta(\vec{v}) < 0} (1 - c_{[\beta]}) l_{[\beta]} &= 0 \end{aligned}$$

therefore $\tilde{R} = \{(1 - c_{[\alpha]}) l_{[\alpha]}, [\alpha] \text{ an equivalence class of roots}\}$ is a finite set of linear functionals whose sum is zero and such that for any codimension 1 hyperplane, the sum of those elements in $\tilde{R}$ lying entirely on a half plane is zero. Therefore $\tilde{R}$ consist of zero linear functionals, so $c_{[\alpha]} = 1$ for all root equivalence classes $[\alpha]$. But this means $l_{\sigma([\alpha])} \circ A = l_{[\alpha]}$. So A is a linear map that sends R_G to $R_{G'}$. $\square$

Remark 3.3.1. *Lemma 3.3.3 implies that the linear part of f , when viewed just as a linear map on $\mathbb{R}^n$, where n is the rank of G (which is also the rank of G') is a scalar multiple of an element $A_f \in O(n)$, where A_f has finite order. (This is because f sends foliations by root kernels of G to that of G' , and elements of R_G (resp. $R_{G'}$) are orthogonal to root kernels.*

The next proposition gives an interpretation to the constant part of f .

Proposition 3.3.1. *Let $\beta \ll \beta' \ll \beta'' \ll 1$ be as in Lemma 3.2.1. Let $S(E_-, E^+, K, h_{\text{bot}}, h_{\text{top}})$ be a generalized $[\alpha]$ slab in B . Suppose $h_{\text{bot}} < z_{\text{bot}} < z_{\text{top}} < h_{\text{top}}$ with $4\beta(h_{\text{top}} - h_{\text{bot}}) \leq (z_{\text{top}} - z_{\text{bot}}) \leq \beta'(h_{\text{top}} - h_{\text{bot}})$, and $|h_{\text{top}} - z_{\text{top}}|, |z_{\text{bot}} - h_{\text{bot}}| > 4\kappa^2\beta''(h_{\text{top}} - h_{\text{bot}})$.*

Then there exists a set $S \subset B$ as in Lemma 3.2.2 such that for all $z \in [z_{\text{bot}}, z_{\text{top}}]$,

$$q(z) = \frac{\mathbf{b}_{[\alpha]}}{\mathbf{b}_{f_*[\alpha]}} z - \frac{1}{\mathbf{b}_{f_*[\alpha]}} \log \frac{|\mathcal{C}_{q(z_{\text{bot}})}g(\pi_{[\alpha]}^+(H) \cap S)|}{|\pi_{[\alpha]}^+(H)|} + O(\hat{\eta} \text{diam}(B)) \quad (12)$$

Remark 3.3.2. In all application of Proposition 3.3.1, we change q by $O(\hat{\eta} \text{diam}(B))$ in order to have equation (12) hold with no error term.

Proof. We know from Lemma 3.3.2 that

$$q(z) = \frac{\mathbf{b}_{[\alpha]}}{\mathbf{b}_{f_*[\alpha]}}(z) + \left(q(z_{\text{bot}}) - \frac{\mathbf{b}_{[\alpha]}}{\mathbf{b}_{f_*[\alpha]}}(z_{\text{bot}}) \right) + O(\hat{\eta} \text{diam}(B))$$

We now find an alternative expression for the term in the bracket.

Let $H_{\text{top}} \subset B$ be a $[\alpha]$ block such that $\alpha_0(H_{\text{top}}) = z_{\text{top}}$. Then, according to equation (3)

$$\log \frac{|\pi_{[\alpha]}^-(H)| e^{-\mathbf{b}_{[\alpha]} z_{\text{top}}}}{|\mathcal{C}_{q(h_1)} g(\pi_{[\alpha]}^-(H))| e^{-\mathbf{b}_{f_*[\alpha]} q(z_{\text{top}})}} = O(\hat{\eta} \text{diam}(B))$$

Simplifying gives

$$\log \frac{|\pi_{[\alpha]}^-(H)|}{|\mathcal{C}_{q(h_1)} g(\pi_{[\alpha]}^-(H))|} = \mathbf{b}_{[\alpha]} z_{\text{top}} - \mathbf{b}_{f_*[\alpha]} q(z_{\text{top}}) + O(\hat{\eta} \text{diam}(B))$$

Together with Corollary 3.2.1 and Lemma 3.3.2 produces

$$\begin{aligned} \frac{1}{\mathbf{b}_{f_*[\alpha]}} \log \frac{|\mathcal{C}_{q(h_2)} g(\pi_{[\alpha]}^+(H) \cap S)|}{|\pi_{[\alpha]}^+(H)|} &= \frac{1}{\mathbf{b}_{f_*[\alpha]}} \log \frac{|\pi_{[\alpha]}^-(H)|}{|\mathcal{C}_{q(h_1)} g(\pi_{[\alpha]}^-(H))|} \\ &= \frac{\mathbf{b}_{[\alpha]}}{\mathbf{b}_{f_*[\alpha]}} z_{\text{top}} - q(z_{\text{top}}) + O(\hat{\eta} \text{diam}(B)) \end{aligned}$$

□

Remark 3.3.3. Now that we know the $\mathbf{A}'$ part of a standard map is affine, we can repeat this entire section for half planes defined for a regular linear functional $\ell \in \mathbf{A}^*$ using the analogous definitions of upper and lower ℓ boundaries given in Section 3.

4 Aligning the linear part of standard maps

Again, we refer to Theorem 3.1. By Lemma 3.3.3, the linear part of f_i 's that appeared in the conclusion of Theorem 3.1 sends R_G of G , to $R_{G'}$ of G' . A priori, the linear parts of the f_i 's do not have to be the same. For a generic G and G' , the group of permutations between R_G and $R_{G'}$ coming from linear maps would be trivial in which case the linear par of f_i 's are identity.

In this section, we show that when rank of G is 2 or higher, the linear parts of the f_i 's have to be the same. When G is rank 1, there are exactly two root classes and the argument follows exactly the same proof as in [EFW2], with the modification of replacing x and y horocycles by left translates of horocycles corresponding to those two root classes.

We aim to prove the following by the end of this section.

Theorem 4.1. *Let G, G' be non-degenerate, unimodular, split abelian-by-abelian Lie groups. Let $\phi : G \rightarrow G'$ be a (κ, C) quasi-isometry. Given $0 < \delta, \eta < \tilde{\eta} < 1$, there exist numbers L_0 such that if $\Sigma \subset \mathbf{A}'$ is a product of intervals of equal size at least L_0 , then there is a subset $U \subset \phi^{-1}(\mathbf{B}(\Sigma)) \subset G$ of relative measure at least $1 - \theta$ such that for any root class horocycle $p V_{[\alpha]} \subset G'$,*

$$d(\phi^{-1}(p V_{[\alpha]} \cap \phi(U)), p' V_{[\beta]} \cap U) = O(\hat{\eta} \text{diam}(\Sigma)), \text{ for some root horocycle } p' V_{[\beta]} \subset G$$

Here $\theta \rightarrow 0$, and $L_0 \rightarrow \infty$ as $\delta, \eta, \tilde{\eta}$ approach zero.

Choose $\delta, \eta \ll 1$. By Lemma ?? and Remark ?? in [P], G is amenable and boxes have small boundary compared to its volume, therefore the same is true of its image under ϕ^{-1} , which means that we can take a sufficiently large box $\mathbf{B}$ and apply Theorem 3.1 to $\phi^{-1}(\mathbf{B})$ to obtain a tiling of $\mathbf{B}$ by images of smaller boxes $\mathbf{B}_i$.

$$\mathbf{B} = \bigsqcup_{i \in \mathbf{I}} \phi(\mathbf{B}_i) \sqcup \Upsilon \quad (13)$$

where $|\Upsilon| \leq \varrho |\mathbf{B}|$, such that there is a subset $\mathbf{I}_0 \subset \mathbf{I}$ of relative measure at least $1 - \varkappa$, where $\varkappa \rightarrow 0$ as $\tilde{\eta} \rightarrow 0$, with the following properties.

For each $i \in \mathbf{I}_0$, there is a subset $U_i \subset \mathbf{B}_i$ of relative measure at least $1 - \theta$, such that $\phi|_{U_i}$ is $\hat{\eta} \text{diam}(\mathbf{B}_i)$ away from a standard map $g_i \times f_i$, where f_i is a affine map respecting root kernels, and whose linear part preserves the set R_G .

Since each U_i has a relative large measure, $U_* = \bigcup_{i \in \mathbf{I}_0} U_i$ relative large measure at least $1 - (\varrho + \theta)$ in $\phi^{-1}(\mathbf{B})$. We write $\mathbf{B}_i = \mathbf{B}(\Omega_i)$ where $\Omega_i \subset \mathbf{A}$ is compact convex, and are all isomorphic to each other.

For a subset $V \subset G'$, we write $\mathbf{I}(V)$ for those $i \in \mathbf{I}$ such that $\mathbf{B}_i \cap V \neq \emptyset$.

Theorem 4.1 finishes the alignment step because of the following consequences.

Corollary 4.0.1. *In the conclusion of Theorem 3.1, the linear part of f_i 's, $i \in \mathbf{I}_0$ are all the same.*

Proof. Let $U_* = \bigcup_{i \in \mathbf{I}_0} U_i \subset G$. Lemma 4.1 implies that for boxes that are within $O(\hat{\eta} \text{diam}(\mathbf{B}))$ of a left translate of $\mathbf{H}$, the f_i 's are the same. Suppose for some $x, y \in \phi^{-1}(\mathbf{B}) \cap U_*$ with $d(\pi_A(x), \pi_A(y)) \geq \hat{\eta} \text{diam}(\mathbf{B})$, the linear part of standard maps supported in neighborhoods of x and y corresponds to distinct permutations $\sigma_{A_x}, \sigma_{A_y}$ from R_G to $R_{G'}$. So there must be a root class $[\alpha]$ such that $[\beta] = \sigma_{A_y}([\alpha])$ is different from $[\Xi] = \sigma_{A_x}([\alpha])$. Let P be a codimension 1 hyperplane such that $\vec{v}_{[\beta]}$ and $\vec{v}_{[\Xi]}$ lie on either side of it.

Without loss of generality, we can take $x V_{[\alpha]}, y V_{[\alpha]}$ such that their intersections with $\mathbf{B}$ have equal measure. Let $x' V_{[\Xi]}$ be a horocycle such that $x' V_{[\Xi]} \cap \mathbf{B}$ is $O(\hat{\eta} \text{diam}(\mathbf{B}))$ Hausdorff distance away from $\phi(x V_{[\alpha]} \cap \phi^{-1}(\mathbf{B}))$, and $y' V_{[\beta]}$ be the corresponding horocycle for $y V_{[\alpha]}$. Since the distance between $\pi_A(x)$ and $\pi_A(y)$ is at least $\hat{\eta} \text{diam}(\mathbf{B})$, we also have $d(\pi_A(x'), \pi_A(y')) > O(\hat{\eta} \text{diam}(\mathbf{B}))$.

Let $\mathcal{F}$ be the set of geodesic segments in direction $P^\perp$ (the unique direction perpendicular to P) with one end point in $x' V_{[\Xi]} \cap \mathbf{B}$ and another in $y' V_{[\beta]} \cap \mathbf{B}$. Then, elements of $\mathcal{F}$ have length at most $O(\text{diam}(\mathbf{B}))$. An element of $\phi^{-1}(\mathcal{F})$ on the other hand, is a path with one end point in $x V_{[\alpha]} \cap \phi^{-1}(\mathbf{B})$ and another in $y V_{[\alpha]} \cap \phi^{-1}(\mathbf{B})$. Since the distance between $\pi_A(x)$ and $\pi_A(y)$ is bigger than $\hat{\eta} \text{diam}(\mathbf{B})$, we can assume without loss of generality that $|\alpha_0(x) - \alpha_0(y)| = \Omega(\tilde{\epsilon} \text{diam}(\mathbf{B}))$, for some $\tilde{\epsilon} \leq \hat{\eta}$ such that $\text{diam}(\mathbf{B}) \ll e^{\tilde{\epsilon} \text{diam}(\mathbf{B})}$. As the measure of $x V_{[\alpha]} \cap \mathbf{B}$ and $y V_{[\alpha]} \cap \mathbf{B}$ are equal, this forces lengths of a large proportion of elements in $\phi^{-1}(\mathcal{F})$ to be at least $e^{\tilde{\epsilon} \text{diam}(\mathbf{B})}$, which is a contradiction because the length of an element of $\phi^{-1}(\mathcal{F})$ is $O(\text{diam}(\mathbf{B}))$. $\square$

Corollary 4.0.2. *Given $0 < \delta, \eta < \tilde{\eta} < 1$, there exist numbers $L_0, \hat{\eta}$ such that if $\Omega \subset \mathbf{A}$ is a product of intervals of equal size at least L_0 , then there is a subset $U \subset \mathbf{B}(\Omega)$ of relative measure at least $1 - \tilde{Q}$, and a standard map $\hat{\phi} = g \times f$ where f is affine defined on it such that*

$$d(\phi|_U, \hat{\phi}) = O(\hat{\eta} \text{diam}(\mathbf{B}(\Omega)))$$

Here $\tilde{Q}$ and $\hat{\eta}$ approach zero as $\tilde{\eta}, \eta, \delta \rightarrow 0$.

Proof. By Theorem 3.1 and Corollary 4.0.1, there is a tiling of $\mathbf{B}(\Omega)$:

$$\mathbf{B}(\Omega) = \bigsqcup_{i \in \mathbf{I}} \mathbf{B}(\omega_i) \sqcup \Upsilon$$

where each ω_i is isometric to $\rho\Omega$ with the properties that there is a subset $\mathbf{I}_0 \subset \mathbf{I}$, with relative measure at least $1 - \varkappa$ such that for $i \in \mathbf{I}_0$, there is a subset $U_i \subset \mathbf{B}(\omega_i)$, with relative measure at least $1 - \theta$, and a standard map $g_i \times f_i$ supported on it such that

$$d(\phi|_{U_i}, g_i \times f_i) = \hat{\eta} \text{diam}(\mathbf{B}(\omega_i))$$

Furthermore, the linear parts of f_i 's are all the same. Denote this common linear map by A_f .

We now basically proceed according to the same steps as in the proof to Theorem 3.1 in [P].

Let $\mathcal{P}^0 = \bigcup_{i \in \mathbf{I}_0} U_i$. Since each U_i has relative measure at least $1 - \theta$ in $\mathcal{P}(\mathbf{B}(\omega_i))$, therefore the measure of $\mathcal{P}^0$ is at least $1 - \theta'$ times that of $\mathcal{P}(\mathbf{B})$, where $\theta' \rightarrow 0$ as $\tilde{\eta}$ approaches zero. Let $\mathcal{L}^0 = \{l \in \mathcal{L}(\mathbf{B}) : |l \cap \mathcal{P}^0| \geq (1 - \theta'^{1/2})|l|\}$. By Chebyshev inequality, the size of $\mathcal{L}^0$ is at least $1 - \theta''$ that of $\mathcal{L}(\mathbf{B})$ where $\theta'' \rightarrow 0$ as $\tilde{\eta}$ approaches zero.

We now show that if $l \in \mathcal{L}^0$, then $\phi(l)$ is close to a geodesic segment.

First we claim that $\pi_A \circ \phi(l)$ is close to a straight line segment in $\mathbf{A}'$. By construction, if $l_i = l \cap \mathcal{P}_i$ for some $i \in \mathbf{I}_0$, then $\pi_A(\phi(l_i))$ is within $\hat{\eta}$ -linear neighborhood of the line segment in direction $A_f \circ l$, and makes an angle at least $\sin^{-1}(\tilde{\eta})$ with root kernels. Since l spend all but $\theta'^{1/2}$ proportion of its length in $\mathcal{P}^0$, we conclude that $\pi_A(\phi(l))$ is within $(\hat{\eta}, \theta'^{1/2}|l|)$ -linear neighborhood of a line segment $\hat{l}$ in direction $A_f \circ l$, and makes an angle at most $\sin^{-1}(\tilde{\eta})$ with root kernels.

Let $h : \phi(l) \rightarrow \hat{l}$ sends each point on $\phi(l)$ to a point on $\hat{l}$ that is closest to its π_A image. We now show that if for some $p, q \in l \cap \mathcal{P}^0$, the line segment connecting $h(\phi(p))$ to $h(\phi(q))$ is orthogonal to $\hat{l}$, then $d(\phi(p), \phi(q)) \leq \hat{\eta}|\phi(l)|$, where $1 \gg \hat{\eta}/2 \gg \theta'^{1/2}$.

Suppose not, then we have two points with above property except that the distance between them is bigger than $\hat{\eta}|\phi(l)|$. Then either $p, q \in \mathcal{P}_i$ for some $i \in \mathbf{I}_0$, or $p \in \mathcal{P}_{\iota_1}, q \in \mathcal{P}_{\iota_2}$ for distinct $\iota_1, \iota_2 \in \mathbf{I}_0$.

The first case implies that $d(\phi(p), \phi(q)) \leq \hat{\eta} \text{diam}(\mathbf{B}_i) < \hat{\eta}|l|$ so this case cannot happen.

So p, q must belong to distinct $\mathbf{B}(\omega_i)$'s. In this case, the lower bound on the distance between $\phi(p)$ and $\phi(q)$ and the knowledge that $\pi_A(\phi(l))$ is within $(\hat{\eta}, \theta'^{1/2}|l|)$ -linear neighborhood of $\hat{l}$ means that there must be two points $x, y \in \mathcal{P}_j \cap l$ for some $j \in \mathbf{I}_0$ such that the oriented line segment connecting $\pi_A(\phi(x))$ and $\pi_A(\phi(y))$ makes an angle of at least $\pi - \sin^{-1}(\tilde{\eta})$ with $\hat{l}$, but this is a contradiction to the Lemma 4.0.1 which says that the linear parts of f_i 's are the same.

Since l spend all but $\theta'^{1/2}$ proportion of its length in $\mathcal{P}^0$, by Lemma ?? in [P], we conclude that $\phi(l)$ is $(\hat{\eta}, \theta'^{1/2}|\phi(l)|)$ weakly monotone, and by Proposition ?? of [P], $\phi(l)$ is within $(\hat{\eta}, \theta'^{1/2}|\phi(l)|)$ -linear neighborhood of a geodesic segment in direction $A_f \circ l$.

So now the ϕ image of each element of $\mathcal{L}^0$ is close to a geodesic segment and we can apply Lemma ?? in [P] to obtain a subset $\mathcal{F}^0 \subset \mathcal{F}(\Omega)$ of relative large measure such that if $\gamma \in \mathcal{F}^0$, $\phi(f)$ is within $\eta \text{diam}(\gamma)$ Hausdorff neighborhood of another flat. We now apply the same argument as in the proof of Theorem 3.1 using quadrilaterals to obtain a standard $\hat{f} \times \hat{g}$ on $\mathcal{P}^0$, where linear part of $\hat{f}$ is A_f . $\square$

4.1 S graph, $\hat{S}$ graph and the H graph

We continue with the setting from equation (13) and discretize $\mathbf{B}$ in this section so that it reflects the structures of the standard maps $\hat{\phi}_i$, for $i \in \mathbf{I}_0$.

The S graph Take $\mathcal{G}'$, a ρ_1 net in G' (the range space) and connect $x, y \in \mathcal{G}'$ by an edge if their π_A images are within ρ_1 of each other and $d(x, y) \leq 10\rho_1$. We metrize this graph by letting lengths of edges be the distance between the corresponding points in G' , so all edges have length $O(\rho_1)$. We refer to the discretization restricted to $\mathbf{B}$ as the S graph.

Recall that each box $\mathbf{B}_i$ tiling the set $\phi^{-1}(\mathbf{B})$ is isometric to $\mathbf{B}(\omega)$, the box associated to some convex compact set $\omega \subset \mathbf{A}$. Let $\rho_i, i = 2, 3, 4, 5$, be numbers such that $\rho_i \ll \rho_{i+1} \ll \text{diam}(\omega)$. Fix $0 < \beta \ll \beta' \ll \beta'' \ll 1$ such that $\hat{\eta} \ll \beta$.

The set $E_\ell(p)$ Let ℓ be a regular linear functional of $\mathbf{A}$ with unit norm. For $p \in G$, we write $E_\ell(p)$ for the left coset $p W_\ell^+ \rtimes \ker(\ell)$. A left coset of $W_\ell^+ \rtimes \ker(\ell)$ is also the pre-image of a point under the projection map $G \rightarrow W_\ell^- \rtimes \mathbb{R}\vec{v}_\ell$.

We say $E_\ell(p)$ is *favourable* if

(i)

$$|E_\ell(p) \cap U_*| \geq \left(1 - \vartheta^{1/2}\right) |E_\ell(p) \cap \phi^{-1}(\mathbf{B})|$$

where ϑ is the relative proportion of U_*^c in $\phi^{-1}(\mathbf{B})$.

(ii) For any point q that is such that $\ell(\pi_A(p)) - \ell(\pi_A(q)) = \rho_5$

$$|E_\ell(q) \cap U_*| \geq \left(1 - \vartheta^{1/2}\right) |E_\ell(q) \cap \phi^{-1}(\mathbf{B})|$$

We say $E_\ell(p)$ is *very favourable* if in addition, for every $i \in \mathbf{I}_0$ such that $|E_\ell(q) \cap U_i| \geq (1 - \theta^{1/3}) |E_\ell(q) \cap \mathbf{B}_i|$, the intersection between $\pi_\ell^-(E_\ell(q) \cap \mathbf{B}_i)$ and $\tilde{E}_{**}$ is non-empty, where $\tilde{E}_{**}$ is the union of E_{**} in those $\mathbf{B}_i$'s, $i \in \mathbf{I}_0 \cap \mathbf{I}(E_\ell(p))$ from Lemma 3.2.1 and Remark 3.3.3. Here θ is an upper bound for U_i^c in $\mathbf{B}_i$.

Since U_* has relative large measure in $\phi^{-1}(\mathbf{B})$, it follows that if we fix a chamber $\mathbf{b}$, there is a left translate of $W_\mathbf{b}^+$, $x_0 W_\mathbf{b}^+$, and a function $\varkappa$ with the following properties:

- A. The measure of $x_0 W_\mathbf{b}^+ \cap U_*$ is at least $1 - \varkappa$ times that of $x_0 W_\mathbf{b}^+$.
- B. There is a subset $\mathcal{S}_0 \subset \mathbf{b}$ of relative measure at least $1 - \varkappa$ such that for every $\ell \in \mathcal{S}_0$, and $p \in U_\mathbf{b}$, $E_\ell(p)$ is very favourable.

where $\varkappa$ approaches zero as our initial data of δ , η and $\tilde{\eta}$ approach zero.

Fix such a left coset of W_b^+ and for every $\ell \in \mathcal{S}_0$, write $\tilde{H}_\ell$ for $E_\ell(x_0) \cap \phi^{-1}(\mathbf{B})$ and H_ℓ for the subset of $Sh(E_\ell(x_0), \rho_5) \cap \phi^{-1}(\mathbf{B})$ whose ℓ value is $\ell(x_0) - \rho_5$.

The sets $\mathbf{I}_g(\ell)$, good and bad ℓ boxes For each $\ell \in \mathcal{S}_0$, let $\mathbf{I}_g(\ell) \subset \mathbf{I}_0$ consisting of indices such that

$$|H_\ell \cap U_i| \cap \left(1 - \vartheta^{1/3}\right) |H_\ell|$$

A box $\mathbf{B}_i$ in the domain is called a *good ℓ box* if $i \in \mathbf{I}_g(\ell)$, and bad otherwise.

Shadows of H_ℓ and $\phi(H_\ell)$ Fix a $\ell \in \mathcal{S}_0$. Let $h_1 = \ell(H_\ell) - \beta \text{diam}(\omega)$ and $h_2 = \ell(H_\ell) - (\beta + \frac{\beta'}{2}) \text{diam}(\omega)$. Since $E_\ell(x_0)$ is very favourable, Lemma 3.2.1 says that with those chosen h_2 and h_1 , for each $i \in \mathbf{I}_g(\ell)$,

$$|Sl_2^1(H_\ell) \cap U_i| \geq (1 - c_2) |Sl_2^1(H_\ell)|$$

For each j between h_2 and h_1 , if we write $\rho(j)$ for the relative proportion of $Sl_2^1(H_\ell) \cap \ell^{-1}(j) \cap U_i^c$ in $Sl_2^1(H_\ell) \cap \ell^{-1}(j)$, the above condition means that

$$\sum_{j=h_2}^{h_1} \rho(j) \leq 2c_2$$

which means that $\rho(h_0^i) \leq 2\sqrt{c_2}$ for some $h_0^i \in [h_2, h_1]$.

The shadow of H_ℓ , denoted by $W(H_\ell)$, is now defined as the union of $\cup_{i \in \mathbf{I}_g(\ell)} Sl_2^1(H_\ell) \cap \ell^{-1}(h_0^i)$ and $\cup_{i \in \mathbf{I}(H_\ell) \setminus \mathbf{I}_g(\ell)} Sl_2^1 \cap \ell^{-1}(h_1)$.

Let $S = U_* \cap S\left(\pi_\ell^-(\tilde{H}_\ell), \pi_\ell^+(\tilde{H}_\ell), \pi_A(\tilde{H}_\ell), \ell(\tilde{H}_\ell), \ell(\tilde{H}_\ell) + \beta'' \text{diam}(\omega)\right)$. For each $i \in \mathbf{I}_g(\ell)$, we define $\hat{W}_i(H_\ell)$ as $\tilde{Sl}_2^1(H_\ell, S) \cap (f_*\ell)^{-1}(q(h_0^i))$. The shadow of $\phi(H_\ell)$, denoted by $\hat{W}(H_\ell)$ is now defined as the union of $\hat{W}_i(H_\ell)$'s, where i ranges over $\mathbf{I}_g(\ell)$.

Shadow vertices We now define a set of ℓ *shadow vertices* in the discretization of $\mathbf{B}$. By shifting the discretization, we can assume that $\hat{W}(H_\ell)$ contains a ρ_1 net of S -vertices. Every S -vertex in $\hat{W}(H)$ is a shadow vertex.

Furthermore, a ℓ shadow vertex is *good* if it lies in $\phi(U_*)$, belonging to ℓ plane containing a point of $\phi(U_*)$, and is at least $10\kappa\beta'' \text{diam}(\omega)$ from $\partial\mathbf{B}$.

If a shadow vertex is not good, then it is bad. We enlarge the set of bad shadow vertices by declaring any S -vertex in the ρ_1 neighborhood of $\phi(U_*^c \cap W(H))$ a bad shadow vertex, even if it was a good shadow vertex by our previous definition.

The bad shadow vertices in $N_{\rho_1} \phi(U_*^c \cap W(H))$ are not necessarily close to $\hat{W}(H)$, even if they come from good boxes[†]. While these bad shadow vertices are not well controlled, they make up a small proportion of all shadow vertices and so do not interfere with the geometric argument in the next section. See Lemma 4.1.1 below.

For either good or bad boxes, the number of ℓ shadow vertices coming from $\mathbf{B}_i$ is proportional to the size of the set $H_\ell \cap \mathbf{B}_i$. The proportionality constant depends only on κ , C , G and G' .

[†]Because it's the bad set in the good box, which we have no control over. Anything termed 'bad' basically comes from sets that we have no control over: everything in bad boxes, and bad sets U_i^c 's in good boxes.

Lemma 4.1.1. *For each $\ell \in \mathcal{S}_0$, there is a constant c_5 depending on our initial data $\delta, \eta, \tilde{\eta}$ such that the proportion of bad shadow vertices is at most c_5 which approach zero as our initial data go to zero's.*

Proof. Bad shadow vertices are defined in two stages. First we have the set S_1 of vertices in $\hat{W}(H_\ell)$ that are either within $10\kappa\beta''diam(\omega)$ of $\partial\mathbf{B}$ or outside of $\beta'diam(\omega)$ neighborhood of a point in $\phi(U_*)$ whose ℓ value is h_0^i smaller than $\ell(H_\ell)$. That this set has small measure in $\hat{W}(H_\ell)$ follows from two facts. First, the subset that are close to $\partial\mathbf{B}$ has relative small measure by Lemma 2.2.1.

Second, if the proportion of S_1 in $\hat{W}(H_\ell)$ is θ , then the set of points in $\tilde{S}l_2^1(H_\ell) \cap \phi(U_*^c)$ contained in a ℓ half plane through a point of S_1 has measure at most θ relative to $\tilde{S}l_2^1(H_\ell)$. However by Lemma 3.2.2, the proportion of $\tilde{S}l_2^1(H_\ell) \cap \phi(U_*^c)$ in $\tilde{S}l_2^1(H_\ell)$ is at most c_4 . Therefore $\theta \leq c_4$.

In the second stage, we enlarge the set of bad vertices in $\hat{W}(H)$ by adding the set $N_{\rho_1}\phi(U_*^c \cap \hat{W}(H_\ell))$. That this set has small measure follows from our choice of h_0^i . $\square$

The $\hat{S}$ -graph We now modify the S -graph near $\phi(\mathbf{H}_\ell)$ so that it reflects divergence property dictated by standard maps.

For $x \in W_\ell^+$, $y \in W_\ell^-$ and $t \in \mathbb{R}$, we write $\gamma_{x,y}(t)$ for the set $(x, y) \ell^{-1}(t)$, and $\gamma_{x,y}([c, d])$ for the $\cup_{t \in [c, d]} \gamma_{x,y}(t)$, which is a ℓ half plane of length $|c - d|$.

Let K_i be the union of $\gamma_{x,y}([q(h_0^i), q(\ell(\tilde{H}_\ell)) - \rho_5/4])$'s that have non-empty intersection with $\hat{W}_i(H_\ell)^\dagger$. We begin by replacing K_i as a subset of the S graph by disjoint union of $\gamma_{x,y}$'s, then define the $\hat{S}$ graph by declaring a new set of vertices and a new incidence relation on K_i .

For each $t_j \in \frac{1}{\rho_1}(q(\ell(\tilde{H}_\ell)) - \rho_5/4 - q(h_0^i))$, call the S vertices of $\gamma_{x,y}(t_j) \in K_i$ *pre-vertices*. Recall that for each linear functional Ξ , we can evaluate $\Xi(p)$ where $p \in G$ to be the Ξ value of $\pi_A(p)$. In the range, we tile left cosets of $W_{(f_i)_*\ell}^-$ in $((f_i)_*\ell)^{-1}(t_j)$ by rectangles T_- 's of diameter $10\rho_1$; in the domain, we tile left cosets of W_ℓ^+ in $\ell^{-1}(q_i^{-1}(t_j))$ by rectangles T_+ 's of diameter $10\kappa^2\rho_1$. We identify two vertices p, q if

(i) p, q are in the same T_- .

(ii) $\phi_i^{-1}(p)$ and $\phi_i^{-1}(q)$ are in the same T_+ which has the property that

$$|\partial_\ell^-(T_+) \cap \tilde{E}_{**}| \geq 1/2 |\partial_\ell^-(T_+)|$$

where $\tilde{E}_{**}$ is the union of E_{**} coming from each $\mathbf{B}_i$, $i \in \mathbf{I}_0$ as in Lemma 3.2.1.

We also remove any edges in K_i that ends at a bad shadow vertex. A $\hat{S}$ vertex is called *regular* unless it arise from the procedure above, in which case it is called *irregular*.

In our original S graph, every point has the same valence provided the vertex is not close to the boundary. However, upon the changes made for the $\hat{S}$ graph, the 'homogeneous'-ness of valence is not so clear. This next lemma says that we only change the valence by bounded amount, so that $\hat{S}$ graph is essentially homogeneous away from boundary.

[†]Note that $\hat{W}_i(H_\ell)$ is only defined for $i \in \mathbf{I}_g(\ell)$.

Lemma 4.1.2. *There exist constants M_l, M_u depending only on κ, C such that for any two $\hat{S}$ vertices v_1, v_2 , the ratio of numbers of $f_*\ell$ half planes through them is bounded between M_l and M_u .*

Proof. Let v be an irregular vertex, and let $z_{top} = \max(f_*\ell)(\mathbf{B})$. In the following calculations, we use $\approx$ to mean that up to a multiplicative error of $e^{O(\hat{\eta} \text{diam}(\omega))}$. First, the number of $f_*\ell$ half planes of length $z_{top} - (f_*\ell)(v)$ containing v and $(f_*\ell)^{-1}(z_{top})$ is $\approx e^{\mathbf{b}_{f_*\ell}(z_{top} - f_*\ell(v))}$.

As v is an irregular vertex, there exists a ℓ block H' in $\mathbf{B}_i$ such that $v \in \phi_i(H')$ and $\partial_\ell^-(H')$ contains a point of E_{**} . The number of $f_*\ell$ half planes containing v and $\hat{W}_i(H_\ell)$ (whose $f_*\ell$ and q_i value is h_0^i) is

$$\begin{aligned}
& \approx |\mathcal{C}_{q_i(h_0^i)} g_i(\pi_\ell^-(H'))| e^{-\mathbf{b}_{f_*\ell} h_0^i} && \text{by (3)} \\
& \approx |\mathcal{C}_{q_i(h_0^i)} g_i(\pi_\ell^-(H'))| \frac{|\mathcal{C}_{q_i(h_2)} g(\pi_\ell^+(H') \cap S)|}{|\pi_\ell^+(H')|} e^{-\mathbf{b}_{\ell q_i^{-1}}(h_0^i)} && \text{by Proposition 3.3.1} \\
& \approx |\pi_\ell^-(H')| e^{-\mathbf{b}_{\ell q_i^{-1}}(h_0^i)} && \text{by 3.2.1} \\
& \approx e^{\mathbf{b}_\ell(\ell(H') - q_i^{-1}(h_0^i))} && \text{by (3)} \\
& = e^{\mathbf{b}_{f_*\ell}(q_i(\ell(H')) - h_0^i)} && \text{by Proposition 3.3.1}
\end{aligned}$$

□

The H graph We now define the H graph as a subgraph of the $\hat{S}$ graph. A good ℓ vertex is an irregular $\hat{S}$ vertex $v \in K_i$ such that $f_*\ell(v)$ is within $O(\rho_1)$ of $q_i(\ell(H_\ell))$ and the W_ℓ^- coordinate of $\phi_i^{-1}(v)$ is within $O(\rho_1)$ of the W_ℓ^- coordinate of H_ℓ . A bad ℓ vertex is a bad ℓ shadow vertices, which is always a regular $\hat{S}$ vertices.

A vertex of the H graph is either good or bad. Good (resp. bad) H vertices is the union over $\ell \in \mathcal{S}_0$, all the good (resp. bad) ℓ vertices. If a vertex comes from the ϕ image of $x_0 W_b^+ \cap U_*$, then it is called a $\mathbf{b}$ vertex.

An edge of the H graph is concatenation of edges in the $\hat{S}$ graph that connects two H vertices, or connect a good H vertex with $\partial\mathbf{B}$. An $\mathcal{E}_0$ edge is one that connects two good H vertices or one good H vertex with $\partial\mathbf{B}$.

4.2 Some geometric lemmas

The idea behind the proof of Theorem 4.1 is the following observation. In H_{n+1} (as in Lemma 2.1.1) suppose two travelers leave a common starting point via segments of diverging vertical geodesics. If they are to meet up again without having to travel for long, then they can only do so in some neighborhood of the starting point.

The projection Π_ℓ and the function $\rho_\ell(\cdot, \cdot)$ Let $\Pi_\ell : G \rightarrow W_\ell^- \rtimes \mathbb{R}\vec{v}_\ell$ be the projection defined by $(\mathbf{x}, \mathbf{t}) \mapsto ([\mathbf{x}]_{W_\ell^-}, \ell(\mathbf{t}))^\dagger$. Let H be a ℓ block, and write $\rho_\ell(p, q) = (\Pi_\ell(p), \Pi_\ell(q))_{\Pi_\ell(H)}$ for the Gromov product of $\Pi_\ell(p)$ and $\Pi_\ell(q)$ with respect to $\Pi_\ell(H)$ in the negatively curved space $W_\ell^- \rtimes \mathbb{R}\vec{v}_\ell$.

$^\dagger[\mathbf{x}]_{W_\ell^-}$ denotes for the W_ℓ^- coordinate of $\mathbf{x}$

Recall that for three points x, y, z in a metric space, the Gromov product is defined as

$$(y|z)_x = \frac{1}{2} (d_X(x, y) + d_X(x, z) - d_X(z, y))$$

In a δ hyperbolic space X , the geodesic joining y to z , γ_{yz} , satisfies

$$d_X(\gamma_{yz}, x) - \delta \leq (y|z)_x \leq d_X(\gamma_{yz}, x)$$

For the remaining of this subsection, H denotes for a ℓ block. Also, until the end of Section 4, ϵ is a positive number less than 1 such that $O(\text{diam}(\mathbf{B})) \ll e^{\epsilon \text{diam}(\mathbf{B})}$, and $\hat{\eta} \text{diam}(\omega) \ll \epsilon \text{diam}(\mathbf{B})$. The length of any path considered in this section is less than $e^{\epsilon \text{diam}(\mathbf{B})}$.

In the next lemma, we list some properties of ρ_ℓ , where $\approx$ is used to denote two quantities whose ratio depends only on κ, C, ℓ and the space G .

Lemma 4.2.1.

(i) Suppose $d(p', p) \ll d(p, H)$, $d(q', q) \ll d(q, H)$, and $\rho_\ell(p, q) \ll \min\{d(p, H), d(q, H)\}$. Then

$$\rho_\ell(p, q) \approx \rho_\ell(p', q')$$

(ii) Suppose $\ell(p') < \ell(p)$, $\ell(q') < \ell(q)$, and each of the pairs (p, p') , (q, q') can be connected by a geodesics having at least η proportion of the component in direction $\vec{v}_\ell$.

If $d(p, H), d(q, H) \gg \rho_\ell(p, q)$, then

$$\rho_\ell(p, q) \approx \rho_\ell(p', q')$$

(iii) If $\rho_\ell(p, q), \rho_\ell(q, q') \gg s$, then $\rho_\ell(p, q') \gg s$.

Lemma 4.2.2. Suppose $p, q \in G$ can be connected by a path $\hat{\gamma}$ of length less than $e^{\epsilon R}$ such that

(i) $\ell(\pi_A(p)), \ell(\pi_A(q)) \leq \ell(H) - \rho_4$

(ii) The ℓ values of $\hat{\gamma}$ decreases for at least ϵR units at both ends, and the ℓ values of points on the the remaining subsegment are no more than $\ell(H) - \epsilon R$.

Then, $\rho_\ell(p, q) \geq \Omega(\rho_4)$.

Proof. Let p' and q' be the closest points to p and q whose ℓ value first dips below $\ell(H) - \epsilon R$. Since the length of $\hat{\gamma}$ is less than $e^{\epsilon R}$, so is the length of $\Pi_\ell(\hat{\gamma})$, which connects $\Pi_\ell(p)$ and $\Pi_\ell(q)$, as well as the subsegment of $\Pi_\ell(\hat{\gamma})$ between $\Pi_\ell(p')$ and $\Pi_\ell(q')$. This means that if $\Pi_\ell(p) \neq \Pi_\ell(q)$, then any path connecting $\Pi_\ell(p')$ to $\Pi_\ell(q')$ whose ℓ value stays ϵR units below $\ell(H)$ would have length at least $e^{\epsilon R}$, contradicting the assumption about the length of $\hat{\gamma}$. $\square$

Lemma 4.2.3. Let p_0, q_0 be good ℓ vertices, and γ be an $\mathcal{E}_0$ edge connecting them. Let $p, q \in \gamma$ be points that are within the same good box as p_0 and q_0 respectively. Then the following holds:

(i) Except near the end points, γ never pass through any irregular ℓ vertices.

(ii) we have $\rho_\ell(\hat{\phi}^{-1}(p), \hat{\phi}^{-1}(q)) \geq \Omega(\rho_4)$

Proof. Starting from p_0 , let p_1 be the first place where γ hits $\hat{W}(H_\ell)$. As γ is a $\mathcal{E}_0$ edge, it doesn't hit a bad ℓ shadow vertex, then there exists a good ℓ shadow vertex $p'_1 \in U_* \cap W(H_\ell)$ such that $\hat{\phi}(p'_1) = p_1$ and $d(\phi^{-1}(p_1), p'_1) = O(\hat{\eta} \text{diam}(\omega))$. Note that p'_1 and $\phi^{-1}(p_1)$ are both $\Omega(\epsilon \text{diam}(\mathbf{B}))$ away from $\partial(\phi^{-1}(\mathbf{B}))$.

Let $p'_2 = \phi^{-1}(p_2)$ be the next point after p'_1 when $\phi^{-1}(\gamma)$ intersects $U_* \cap W(H_\ell)$. We know there must be such a point because the length of γ is less than $e^{\epsilon \text{diam}(\mathbf{B})}$, so whenever it moves transverse to H_ℓ , it must do so in $\ell^{-1}[h_0^i, \infty]$.

Since γ does not hit a bad shadow vertex, $p'_2 \in \ell^{-1}(h_0^i)$ at $i \in \mathbf{I}_g(\ell)$, so p_2 is a good shadow vertex. Therefore continuing γ after p_2 hits a vertex in $\phi(U_*) \cap H_\ell$, which is a good ℓ vertex, and must be q_0 .

By Lemma 4.2.1

$$\rho_\ell(\hat{\phi}^{-1}(p), \hat{\phi}^{-1}(q)) = \rho_\ell(\hat{\phi}^{-1}(p_1), \hat{\phi}^{-1}(p_2)) \approx \rho_\ell(\phi^{-1}(p_1), \phi^{-1}(p_2)) \geq \Omega(\rho_4)$$

□

Lemma 4.2.4. *Suppose p_0, q_0 are good ℓ vertices and $\overline{p_0 q_0}$ is a $\mathcal{E}_0$ edge connecting them. If for some $\rho_1 \ll s \ll \rho_3 \ll \rho_4$, we have $p, q \in \overline{p_0 q_0}$ in the same good box as p_0 and q_0 satisfying*

$$(f_i)_* \ell(p) - (f_i)_* \ell(p_0) = (f_j)_* \ell(q) - (f_j)_* \ell(q_0) = s, \text{ where } i, j \in \mathbf{I}_g(\ell)$$

Then, there is a ℓ block H'_ℓ such that p, q are within $O(\rho_1)$ of $\hat{\phi}(H'_\ell)$.

Proof. Let p', q' be points on $\overline{p_0 q_0}$ close to where it enters respective good boxes. By Lemma 4.2.3, $\rho_\ell(\hat{\phi}^{-1}(p'), \hat{\phi}^{-1}(q')) \geq \Omega(\rho_4)$. Since $s \ll \rho_3 < \rho_4$, by Remark 3.3.1, we conclude that the ℓ values of $\hat{\phi}_i^{-1}(p)$ and $\hat{\phi}_j^{-1}(q)$ are the same, both are s lower than that of H_ℓ . Since $s < \rho_3 \ll \rho_4$, $\Pi_\ell(\hat{\phi}_i^{-1}(p))$ and $\Pi_\ell(\hat{\phi}_j^{-1}(q))$ are the same point, which is the same as saying that $\hat{\phi}_i^{-1}(p)$ and $\hat{\phi}_j^{-1}(q)$ are within $O(\rho_1)$ of a ℓ block H'_ℓ as claimed. □

Lemma 4.2.5. *Let n be 4 or 6. Suppose for $0 \leq i \leq n-1$, p_i are $\hat{S}$ vertices whose ϕ pre-images support standard maps. Let $\overline{p_{i-1} p_i}$ be subsegments of $\mathcal{E}_0$ edges in H graph, where the indices are counted mod n .*

Let $r(p_i) = \min\{|(f_i)_ \ell(v) - (f_i)_* \ell(p_i)|, v \text{ is a good } \ell \text{ vertex}\}$, where $\phi_i = g_i \times f_i$ is the standard map supported in a neighborhood of $\phi^{-1}(p_i)$.*

Suppose there is an index k such that $r(p_k) \ll \rho_4$, and for all $i \neq k$, $r(p_i) > r(p_k) + 2\rho_1$. Then $\overline{p_k p_{k+1}}$ and $\overline{p_k p_{k-1}}$ cannot have only p_k in common.

Proof. we can assume $k = 0$. Let H'_ℓ be the ℓ block passing through $\hat{\phi}^{-1}(p_0)$. By Lemma 4.2.4, we can consider H'_ℓ in place of H_ℓ . Namely, we can replace the appearance of H_ℓ in the definition of ℓ vertices by H'_ℓ . Let p_i^+ and p_i^- be the first and last time that $\overline{p_{i-1} p_i}$ leaves $\hat{W}(H_\ell)$.

By Lemma 4.2.3, for all $i \neq 0$, $\rho_\ell(\hat{\phi}^{-1}(p_{i-1}^+), \hat{\phi}^{-1}(p_i^-)) \geq \Omega(\rho_4)$, and $\rho_\ell(\hat{\phi}^{-1}(p_0^-), \hat{\phi}^{-1}(p_0^+)) \leq \rho_1$. This is a contradiction to Lemma 4.2.1 □

4.3 Proof of Theorem 4.1

Let $\mathbb{R}^k \rtimes_\tau \mathbb{R}$ be a rank 1 space with roots $\alpha_i, -\beta_j$'s, where $\alpha_i, \beta_j > 0$, and $\tau(t)$ is matrix consisting of blocks of the form $e^{\alpha_i t} N_i(\alpha_i t)$, $e^{-\beta_j t} N_j(\beta_j t)$ where N_i, N_j are unipotent matrices with polynomial

entries. For the following two lemmas, let $B[T]$ denote for the subset $\left(\prod_i \Omega_i \times \prod_j U_j\right) \rtimes [-T, T]$, where $\Omega_i \subset V^{\alpha_i}$, and $U_j \subset V^{-\beta_j}$, and $\alpha_i, \beta_j > 0$, $|\Omega_i|e^{-\alpha_i T}, |U_j|e^{-\beta_j T} \geq 1$. We call the set

$$\left(\prod_i \Omega_i \times \prod_j U_j\right) \rtimes -T \cup \left(\prod_i \Omega_i \times \prod_j U_j\right) \rtimes T$$

the *top and bottom* of $B[T]$, denoted by $\bar{\partial}B[T]$.

Lemma 4.3.1. *The total number of geodesics in $B[T]$ is $e^{T(\sum_i \alpha_i + \sum_j \beta_j)} \prod_{i,j} |\Omega_i| |U_j|$, and the number of geodesics in $B[T]$ through each vertex is $e^{T(\sum_i \alpha_i + \sum_j \beta_j)}$.*

Proof. To specify a geodesic in $B[T]$, we need to specify its coordinates in α_i, β_j root spaces, and for every choice of α_i and β_j coordinate, there is a unique geodesic segment in $B[T]$ going from the top to the bottom. The number of different coordinates in α_i root spaces is $\prod_i |\Omega_i|e^{\alpha_i T}$, and those in β_j root spaces is $\prod_j |U_j|e^{\beta_j T}$, so the number of geodesics is

$$\left(\prod_i |\Omega_i|e^{\alpha_i T}\right) \left(\prod_j |U_j|e^{\beta_j T}\right)$$

We know that in $B[T]$

no. of geodesics $\times$ no. of vertices on a geodesic = no. vertices $\times$ no. of geodesics through each vertex

The number of vertices on each geodesic is $2T$, and the number of vertices is $2T(\prod_i |\Omega_i|)(\prod_j |U_j|)$, and we now see the number of geodesics through each point is indeed as claimed. $\square$

Lemma 4.3.2. *Hypothesis as in Lemma 4.3.1. Now further assume that $\mathbb{R}^k \rtimes_{\tau} \mathbb{R}$ is unimodular, and let the common values of $\sum_i \alpha_i$ and $\sum_j \beta_j$ be m . Let $X \subset B[T]$ be a subset of vertices. If $\mathcal{F}$ is a family of geodesics in $B[T]$ with size $\sigma e^{2mT} \prod_{i,j} |\Omega_i| |U_j|$, where $\sigma \gg \frac{2T}{e^{m\rho_2}}$, then there is a vertex $v \in X$, and two geodesics through v in the same direction that stay together for shorter than ρ_2 units.*

Proof. Suppose the claim is not true. Then for each X vertex v , every pairs of geodesics through v stay together at least ρ_2 units. If v is within ρ_2 neighborhood of top and bottom of $B[T]$, the number of geodesics through v with properties is $\frac{e^{(2T-h(v))m}}{e^{m\rho_2}}$, where h is the height function on $\mathbb{R}^m \rtimes \mathbb{R}$. On the other hand, if v is outside of ρ_2 neighborhood of top and bottom of $B[T]$, the number of geodesics through v with this property is

$$\frac{e^{h(v)\sum_i \alpha_i}}{e^{m\rho_2}} \frac{e^{(2T-h(v))\sum_j \beta_j}}{e^{m\rho_2}} = \frac{e^{2mT}}{e^{2m\rho_2}}$$

The number of X vertices outside of ρ_2 neighborhood of the top and bottom of $B[T]$ is at most $2(T - \rho_2) \prod_{i,j} |\Omega_i| |U_j|$, and the number of X vertices within ρ_2 neighborhood of top and bottom of $B[T]$ is at most $2\rho_2 \prod_{i,j} |\Omega_i| |U_j|$. So the number of geodesics satisfying this scenario is at most

$$\sum_{v \in N_{\rho_2} \bar{\partial}B[T]} \frac{e^{(2T-h(v))m}}{e^{m\rho_2}} + \frac{e^{2mT}}{e^{2m\rho_2}} 2(T - \rho_2) \prod_{i,j} |\Omega_i| |U_j| \leq \frac{e^{2Tm}}{e^{m\rho_2}} (2T) \prod_{i,j} |\Omega_i| |U_j|$$

Since the size of $\mathcal{F}$ is larger than this number, it is not possible that every pairs of geodesics in $\mathcal{F}$ satisfy the scenario described above. So there must be a X vertex v , and two geodesics in $\mathcal{F}$ that stay together for less than ρ_2 units after passing through v . $\square$

In the remaining of this subsection, given a regular vector $\vec{u}$, we write $I_{\lambda, \vec{u}}(p)$, $p \in G'$, for the subset of $pW_{\vec{u}}^+$ that can be reached by two geodesics of length λ in direction $\vec{u}$ in the rank 1 space $G'_{\vec{u}} = \mathbf{H}' \rtimes \mathbb{R}\vec{u}$ containing p , and $I'_{\lambda, \vec{u}}(p)$ for the subset of the left $W_{\vec{u}}^-$ coset that can be reached from p by a geodesic in direction $\vec{u}$ (or in direction $-\vec{u}$ as viewed from p) of length λ .

For the next two propositions, we make the following assumptions.

- (i) Let v be a $\mathfrak{b}$ vertex such that $\phi^{-1}(v)$ locally supports a standard map $\phi_v = f_v \rtimes g_v$. Since f_v is affine and permutes root classes of G to root classes of G' , its linear part induces a permutation from the chambers of G to chambers of G' and we write $(f_v)_*$ for this permutation.
- (ii) Suppose for $\ell \in (\mathbf{A})^*$ a regular linear functional, and $\vec{u} \in \mathbf{A}'$, the vectors $\vec{v}_\ell$ and $(f_v)_*^{-1}\vec{u}$ lie in a common chamber $\mathfrak{b}$ of G .

Proposition 4.3.1. *If λ is a number such that at least σ fraction of geodesics leaving v in direction $\vec{u}$ are unobstructed by H_ℓ vertices for length at least $\lambda + \rho_2$, where $\sigma \gg 2\lambda/e^{\mathbf{b}_{\vec{u}}\rho_2}$. Suppose $\sigma > O(\eta)$ for some $\eta < 1$, then at least $1 - O(\eta)$ fraction of the $\hat{S}$ vertices in $I_{\lambda, \vec{u}}(v)$ are H_ℓ vertices.*

Proof. Let E denote the set of geodesics leaving v in direction $\vec{u}$ that are unobstructed by ℓ vertices of length at least $\lambda + \rho_2$. Let E_λ be the subset of $I'_{\lambda, \vec{u}}(v)$ passing through an element of E . By assumption, we have

$$|E_\lambda| \geq \sigma e^{\mathbf{b}_{\vec{u}}\lambda}$$

Let $\mathcal{F}'_0$ be the union of geodesics leaving E_λ in direction $-\vec{u}$. (as viewed from E_λ). Applying Lemma 4.3.2 to $\mathcal{F}'_0$, where X means ℓ vertices, we see that either there is a ℓ vertex whose $\vec{u}^* = \langle \vec{u}, \pi_A(\cdot) \rangle$ value is at most ρ_2 from that of v , or that there is a ℓ vertex w whose $\vec{u}^*$ value differ from that of v by more than ρ_2 , and two geodesics in $G_{\vec{u}}$ through w_1 that stay together for less than ρ_2 units after passing through w_1 . Suppose the latter happens. Let $x, y \in I'_{\lambda, \vec{u}}(v)$ be two upper end points of those two geodesics, and w_1 be the first time that $\overline{wx}$ diverge from $\overline{wy}$. See Figure 1 below.

Let z be the first time that $\overline{vx}$ diverge from $\overline{vy}$. Applying Lemma 4.2.5 to the loop $z - y - w_1 - x - z$ creates a contradiction. So there is a ℓ vertex whose $\vec{u}^* = \langle \vec{u}, \pi_A(\cdot) \rangle$ value is at most ρ_2 from that of v . That is, there is a ℓ vertex in the ρ_2 neighborhood of $I_{\lambda, \vec{u}}(v)$.

Let $U' \subset I_{\lambda, \vec{u}}(v)$ be those vertices that can be reached by two elements of $\mathcal{F}'_0$. Since every vertex in $I_{\lambda, \vec{u}}(v)$ can be reached by at most $|E_\lambda|$ geodesics in $I'_{\lambda, \vec{u}}(v)$, it follows that the relative measure of U' in $I_{\lambda, \vec{u}}(v)$ is at least $1 - O(\eta)$.

Now suppose $w \in U'$, and let $x, y \in I'_{\lambda, \vec{u}}(v)$ be such that $\overline{wx}, \overline{wy}$ are element of $\mathcal{F}'_0$ that are not obstructed by ℓ vertices. Applying Lemma 4.2.5 to the loop $v - x - w - y - v$, and noting that $r(v) = 0$, $r(x), r(y) \geq \rho_2$, it follows that $r(w) = 0$ (otherwise $r(v)$ would be the smallest, a contradiction), which means that w is a ℓ vertex. $\square$

Proposition 4.3.2. *(Hypothesis as in Proposition 4.3.1) Let $\mathcal{F}$ be the union of geodesics in direction $\vec{u}$ leaving a point of $I_{\lambda, \vec{u}}(v)$. Then at least $1 - O(\eta)$ fraction of $\mathcal{F}$ are unobstructed by H_ℓ vertices for length $\lambda + \rho_2$.*




Figure 1: The loop in the proof of Proposition 4.3.1. Filled boxes are H_ℓ vertices.

Proof. Let $E_\lambda \subset I'_{\lambda, \vec{u}}(v)$, $U' \subset I_{\lambda, \vec{u}}(v)$ and $\mathcal{F}'_0$ be as in Proposition 4.3.1. Note that the measure of U' is at least $1 - O(\eta)$ that of $I_{\lambda, \vec{u}}(v)$.

Let $\mathcal{F}''$ be the set of geodesics leaving U' in direction $\vec{u}$. Let $\mathcal{F}''_{long}$ be the set of geodesics coming from extending elements of $\mathcal{F}''$ by extending ρ_4 units on the $I_{\lambda, \vec{u}}(v)$ [†]. Let H'_ℓ be the ℓ block whose ℓ value is ρ_4 less than $\ell(H_\ell)$. We call the resulting vertices ℓ' vertices if we replace occurrence of H_ℓ in the definition of ℓ vertices by H'_ℓ .

If all elements of $\mathcal{F}''_{long}$ are unobstructed by images of ℓ' vertices, then almost all elements of $\mathcal{F}$ are unobstructed by ℓ vertices, where ‘almost’ here means with relative proportion at least $1 - O(\eta)$. Let U'_{long} be the set of ℓ' vertices that are within ρ_4 of U' .

We have that $|\mathcal{F}''_{long}| \geq (1 - O(\eta))e^{\mathbf{b}_{\vec{u}}(2\lambda + \rho_4)}$. Applying Lemma 4.3.2 allows us to conclude that either there is a ℓ vertex whose $\vec{u}^*$ value is within ρ_2 to $\partial \mathbf{B} \cap G'_{\vec{u}}$, or that there is a ℓ vertex q whose $\vec{u}^*$ value differ from that of $\partial \mathbf{B} \cap G'_{\vec{u}}$ by more than ρ_2 units, and two elements of $\mathcal{F}''_{long}$ that stay together for less than ρ_2 units after passing through q .

Suppose the latter scenario happens. Then there are $w_1, w_2 \in I_{\lambda, \vec{u}}(v)$ and a ℓ' vertex q such that $\overline{w_1 q}, \overline{w_2 q} \in \mathcal{F}''_{long}$. Let q_* be the first point where $\overline{w_1 q}$ and $\overline{w_2 q}$ come together. Then by assumption, the $d(q, q_*) < \rho_2$. Let $x_1 \in I'_{\lambda, \vec{u}}(v)$ be the first point where geodesics in direction $\vec{u}$ leaving w_1 and v first meet, and let $x_2 \in I'_{\lambda, \vec{u}}(v)$ be similarly defined for w_2 and v . Let $r(\cdot)$ now denotes for the distance to the closest ℓ' vertex. Then in the loop $v - x_1 - w_1 - q_* - w_2 - x_2 - v$, (see Figure 2) the r value of all points but q_* are at least ρ_4 , while $r(q_*) \leq \rho_2$, which is a contradiction by Lemma 4.2.5.

[†]Ideally, we would like to apply Lemma 4.3.2 to the family $\mathcal{F}''$ in a rank 1 box of size λ but in order to use illegal circuit we need the stub - vertical segment from H_ℓ instead of being on H_ℓ . Hence the choice of H'_ℓ lower down.




Figure 2: The loop in the Proof of Proposition 4.3.2. Filled boxes are $H_{\ell'}$ vertices.

Therefore if elements of $\mathcal{F}_{long}''$ is to contain a ℓ' vertex, this vertex is within ρ_2 neighborhood of $\partial\mathbf{B} \cap G_{\vec{u}}'$, which is just saying that no elements of $\mathcal{F}_{long}''$ are obstructed by ℓ' vertices, therefore at least $1 - O(\eta)$ proportion of elements in $\mathcal{F}$ are unobstructed by ℓ vertices by construction. $\square$

Theorem 4.2. *Let v be a $\mathfrak{b}$ vertex and $\hat{\phi}_v = f_v \times g_v$ be a standard map supported in a neighborhood of $\phi^{-1}(v)$. Let $\lambda_0 = d(v, \partial\mathbf{B})$. Then at least $1 - O(\eta)$ proportion of vertices in $vW_{(f_v)_*\mathfrak{b}}^+$ are $\mathfrak{b}$ vertices, where $O(\eta)$ is the proportion of geodesics leaving $\phi^{-1}(v)$ that admits a geodesic approximation of length $\Omega(\text{diam}(\omega))$.*

Proof. Let $s_{\vec{u}}$ be the difference in $v_{\vec{u}}^*$ values between $\partial\mathbf{B} \cap G_{\vec{u}}'$ and v .

For a $\rho_2 = \hat{\eta}s_{\vec{u}}$. For every $\ell \in \mathcal{S}_0$, and $w \in I_{\lambda, \vec{u}}(v)$, we let $f_{\ell}(w, \lambda)$ denotes the proportion of geodesics leaving w that are unobstructed by H_{ℓ} vertices for length at least $\lambda + \rho_2$. Let

$$f_{\ell}^*(v, \lambda) = \sup_{w \in I_{\lambda, \vec{u}}(v)} f_{\ell}(w, \lambda)$$

In view of Proposition 4.3.1 and 4.3.2, if $f_{\ell}^*(v, \lambda) \geq O(\eta)$, then $f_{\ell}^*(v, \lambda) \geq 1 - O(\eta)$.

Then, either for all $\lambda \leq s_{\vec{u}}$, we have $f_{\ell}^* \geq 1 - O(\eta)$; or that there is a maximal number $\lambda_{\vec{u}, \ell} - 1$ such that $f_{\ell}^*(v, \lambda_{\vec{u}, \ell} - 1) \geq 1 - O(\eta)$, but $f_{\ell}^*(v, \lambda_{\vec{u}, \ell}) < O(\eta)$. We are done if the latter does not happen. We now proceed to show that this is indeed the case.

In the second scenario, we know that $\lambda_{\vec{u}, \ell} \geq \hat{\eta}\text{diam}(\omega)$, and at least $1 - O(\eta)$ proportion of vertices in $I_{\lambda_{\vec{u}, \ell}, \vec{u}}(v)$ and $I'_{\lambda_{\vec{u}, \ell}, \vec{u}}(v)$ are H_{ℓ} vertices. That is, they are ϕ images of $U_* \cap (x_0 W_{\mathfrak{b}}^+ \rtimes \ker(\ell))$.

Claim: If $\lambda_{\vec{u},\xi} > \lambda_{\vec{u},\ell}$ for $\xi, \ell \in \mathcal{S}_0$, then $\lambda_{\vec{u},\xi} - \lambda_{\vec{u},\ell} > O(\hat{\eta} \text{diam}(\omega))$.

Suppose not. Then we will have subsets, one in $\ker(\xi)$ and one in $\ker(\ell)$ that are within $O(\hat{\eta} \text{diam}(\omega))$ Hausdorff distance from each other. This can only happen if the subsets are within $O(\hat{\eta} \text{diam}(\omega))$ of $\ker(\xi) \cap \ker(\ell)$. But this would mean that most of $I'_{\lambda_{\vec{u},\ell}, \vec{u}}(v)$ come from ϕ images of $x_0 W_{\mathfrak{b}}^+ \rtimes (\ker(\xi) \cap \ker(\ell))$, contradicting the assumption that $\lambda_{\vec{u},\xi}$ is the minimal height t where most of the $I'_{t, \vec{u}}(v)$ are obstructed by H_{ξ} vertices.

In this way, the image of the map $\mathcal{S}_0 \rightarrow [0, s_{\vec{u}}]$ defined by sending $\xi \rightarrow \lambda_{\vec{u},\xi}$ is a $O(\hat{\eta} \text{diam}(\omega))$ discrete set. Let $\hat{\lambda}_{\vec{u}}$ be the minimal image value whose pre-images has positive measure. This means that most elements of $I'_{\hat{\lambda}_{\vec{u}}, \vec{u}}(v)$ and $I_{\hat{\lambda}_{\vec{u}}, \vec{u}}(v)$ are ϕ images of $U_* \cap x_0 W_{\mathfrak{b}}^+$, thus the subset of $\mathcal{S}_0$ consisting of elements ξ such that $\lambda_{\vec{u},\xi} > \hat{\lambda}_{\vec{u}}$ is empty. Since for all $t < \hat{\lambda}_{\vec{u}}$, the pre-images of t in $\mathcal{S}_0$ has zero measure, this means not only does the pre-images of $\hat{\lambda}_{\vec{u}}$ has positive measure, it has full measure.

Now pick another direction $\vec{u}'$ in the same chamber as $\vec{u}$, but not in the $\hat{\eta}^{1/2}$ neighborhood of the $\vec{u}$ orbit under the finite group of affine maps permuting R_g to $R_{G'}$, and repeat the same argument as above to obtain a number $\hat{\lambda}_{\vec{u}'}$ such that most of $I'_{\hat{\lambda}_{\vec{u}'}, \vec{u}'}(v)$ come from ϕ images of $x_0 W_{\mathfrak{b}}^+$.

Pick $y_0 \in I'_{\hat{\lambda}_{\vec{u}}}(v)$ and $y \in I'_{\hat{\lambda}_{\vec{u}'}}(v)$ so that each locally supports a standard map. Take two geodesics leaving y_0 in direction $\vec{u}$ (as viewed from v) that stay together for $t_{\vec{u}}$ units (where $\hat{\eta}^{1/2} \text{diam}(\omega) \ll t_{\vec{u}} \ll \text{diam}(\omega)$) after they leave y_0 , followed by a short segment $\text{diam}(\omega)$ away from v , before joining the geodesics connecting v to y . Let's say they stay together for $t_{\vec{u}'}$ units before coming to a stop at y . See Figure 3

As most of $I'_{\hat{\lambda}_{\vec{u}}}(v)$ and $I'_{\hat{\lambda}_{\vec{u}'}}(v)$ come from $U_* \cap W_{\mathfrak{b}}^+$, for a full measure of $\ell \in \mathfrak{b}$, Lemma 4.2.5 requires us to have $\Pi_{\ell} \phi_{y_0}^{-1}(t_{\vec{u}} \vec{u}) \geq \Pi_{\ell} \phi_y^{-1}(t_{\vec{u}'} \vec{u}')$, as well as $\Pi_{\ell} \phi_{y_0}^{-1}(t_{\vec{u}} \vec{u}) \leq \Pi_{\ell} \phi_y^{-1}(t_{\vec{u}'} \vec{u}')$. This means that $f_{y_0}^{-1}(t_{\vec{u}} \vec{u}) = f_y^{-1}(t_{\vec{u}'} \vec{u}')$, where f_{y_0} and f_y are linear part of standard maps ϕ_{y_0} and ϕ_y . That is, $t_{\vec{u}}/t_{\vec{u}'} \in [1/(1 - \hat{\eta}), 1 + \hat{\eta}]$, and $t_{\vec{u}} \vec{u}$ lies in the $\hat{\eta} \text{diam}(\omega)$ neighborhood of the orbit of $t_{\vec{u}'} \vec{u}'$ under the finite group of linear maps that permutes R_G to $R_{G'}$. But this contradicts our choice of $\vec{u}'$ and $t_{\vec{u}}$. $\square$

Proof. of Theorem 4.1 As any root class horocycle is the intersection of finitely many left translates of $W_{\mathfrak{b}}^+$, where $\mathfrak{b}$ is a chamber, it suffices to show that the analogue claim holds for left translates of $W_{\mathfrak{b}}^+$ of G' in place of left translates of $V_{[\alpha]}$ of G' .

We start with $\mathbf{B}(\Sigma)$ sufficiently large so that we can apply Theorem 3.1 to $\phi^{-1}(\mathbf{B}(\Sigma))$ and obtaining a tiling as in equation (13). Let $U_* = \bigcup_{i \in \mathbf{I}_0} U_i$. Since U_* has large measure relative to $\phi^{-1}(\mathbf{B}(\Sigma))$, for a fixed chamber $\mathfrak{b}$, we can find a large subset $U_{\mathfrak{b}} \subset U_*$ with the property that for every point $p \in U_{\mathfrak{b}}$, there is a subset $\mathcal{S}_0 \subset \mathfrak{b}$ of relative proportion at least $1 - \vartheta$ such that $E_{\ell}(p)$ is very favourable for every $\ell \in \mathcal{S}_0$. By constructing the corresponding $\hat{S}$ and H graph, application of Theorem 4.2 to a point $v \in \phi(U_{\mathfrak{b}})$ shows that the ϕ^{-1} image of $v W_{\mathfrak{c}}^+ \cap \mathbf{B}(\Sigma)$ is $O(\hat{\eta} \text{diam}(\mathbf{B}(\Sigma)))$ away from a left translate of $W_{\mathfrak{b}}^+$, where $\mathfrak{c}$ is the image of $\mathfrak{b}$ under the linear part of the standard map supported in a neighborhood of $\phi^{-1}(v)$. $\square$




Figure 3: The loop that prevents blocking. Filled boxes represent $\mathfrak{b}$ vertices. The left hand is the π_A projection image of the loop on the right.

5 Patching

In the previous section, we aligned the linear part of standard maps appeared in Theorem 3.1 by showing that they are all the same. In this last section, we remove the condition that standard maps are only defined for a subset of relative large measure and align the translational part of the standard maps by adopting the procedure used to achieve this in [EFW1].

5.1 A weak version of an affine map

We have by now seen that given a box, there is a subset of large measure supporting a standard map. In this subsection, by controlling the sizes of increasingly larger and larger boxes, we get rid of the 'subset of large relative measure' and extend the result to all pairs of points $p, q \in G$ on the same flat. The precise statement is the following:

Theorem 5.1. *Let G, G' be non-degenerate, unimodular, split abelian-by-abelian Lie groups and $\phi : G \rightarrow G'$ be a (κ, C) quasi-isometry between them. Given $0 < \delta, \eta \ll \tilde{\eta} < 1$, then there exists $\tau < 1$, M depending on $\delta, \eta, \tilde{\eta}$ and QI constants of ϕ such that whenever x, y belong to the same left coset of $\mathbf{H}$,*

$$|\pi_A(\phi(x)) - \pi_A(\phi(y))| \leq \tau d(x, y) + M \quad (14)$$

where $\tau \rightarrow 0$, $M \rightarrow \infty$ as the input parameters approach zeros.

The setup to the proof of theorem 5.1 follows the same sequence of steps as the analogue result in Section 6.1 of [EFW1].

Fix $0 < \delta, \eta \ll \tilde{\eta} < 1$. Let $\Omega \subset \mathbf{A}$ be a product of intervals of size L_0 with barycenter located at the origin of $\mathbf{A}$. By Corollary 4.0.2, there is a subset $\mathcal{P}^0 \subset \mathcal{P}(\Omega)$ of relative large measure which is the support of a standard map $g \times f$ where f is affine. Let $\vartheta \ll 1$ satisfies $|\mathcal{P}^0| \geq (1 - \vartheta)|\mathcal{P}(\Omega)|$, and set $\varrho = \sqrt{\vartheta}$. Let $\mathbf{P}$ be a left translate of $\mathbf{H}$, and we can assume $\mathbf{P}$ contains the identity element. Let $R(\Omega) = \mathbf{B}(\Omega) \cap \mathbf{P}$, and $\mathbf{R}(\Omega) = \bigcup_{g \in R(\Omega)} g(\varrho\Omega)$. The following is a rehash of Corollary 4.0.2.

Corollary 5.1.1. *There is a standard map $f \times g$ where f is affine with linear part A_f , defined on $\mathcal{P}^0 \subset \mathcal{P}(\mathbf{R}(\Omega))$, with $|\mathcal{P}^0| \geq (1 - \varrho)|\mathcal{P}(\mathbf{R}(\Omega))|$, such that $d(\phi|_{\mathcal{P}^0}, f \times g) \leq \hat{\eta} \text{diam}(\mathbf{B}(\Omega))$.*

Furthermore, if $p \in \mathcal{P}^0$, there is a subset $L^0(p) \subset L(p)$ of relative large measure such that the ϕ image of every element $\zeta \in L^0(p)$ is within $\hat{\eta}$ -linear neighborhood of a geodesic segment in direction that of $A_f \circ \zeta$, whose direction makes an angle of at most $\sin^{-1}(\tilde{\eta})$ with root kernels.

The tiling. Choose $\hat{\eta} \ll \varsigma \ll 1$. For each $j \in \mathbb{N}$, set $\Omega_j = (1 + \varsigma)^j \Omega$. We tile $\mathbf{P}$ by $R(\Omega_j)$, where each tile is denoted by $R_{j,\iota}$, $\iota \in \mathbb{N}$. For $p \in G$, we write $R_j[x]$ for the tile in the j -th tiling containing the point in $\mathbf{P}$ that lies on the same flat as x .

Warning. Despite the fact that $\Omega_{j+1} = (1 + \varsigma)\Omega_j$, the number $R_{j,k}$'s needed to cover a rectangle $R_{j+1}[p]$ is very large, on the order of $e^{\sum_{i=1}^{|\Delta'|} \varsigma \max\{\alpha_i(\Omega)\}}$.

The sets U_j For each tile $R_{j,k}$ in the j -th tiling of $\mathbf{P}$, Corollary 5.1.1 produces a subset $\mathcal{P}_{j,k}^0 \subset \mathcal{P}(\mathbf{R}_{j,k})$ of relative large measure. We set

$$U_j = \bigcup_{k \in \mathbb{N}} \mathcal{P}_{j,k}^0$$

In view of Corollary 5.1.1, for any $x \in U_j$,

$$\sup_{y \in \mathbf{R}[x] \cap U_j} |\pi_A(\phi(x)) - \pi_A(\phi(y))| \leq \hat{\eta} \text{diam}(\mathbf{B}(\Omega_j)) \quad (15)$$

Recall that Δ' is the set of roots in G' , and we write n for the rank of G (which is also the rank of G'). We also have the following generalization:

Lemma 5.1.1. *For any $x \in U_j$, and $y \in \mathbf{R}_{j+1}[x] \cap U_j$,*

$$|\pi_A(\phi(x)) - \pi_A(\phi(y))| \leq (|\Delta'|)(n)4\hat{\eta} \text{diam}(\mathbf{B}(\Omega_j))$$

Proof. WLOG, we can assume that there is a horocycle $H_{[\alpha]}$, $[\alpha]$ an equivalence class of roots, that intersect both $\mathbf{R}[x] \cap U_j$ and $\mathbf{R}[y] \cap U_j$. (If not, then we can find a sequence of points $x = p_0, p_1, \dots, p_l = y$ where $l \leq |\Delta'|$ such that for each pair of consecutive points, there is a horocycle intersecting $\mathbf{R}[p_\iota] \cap U_j$ and $\mathbf{R}[p_{\iota+1}] \cap U_j$). Let $x_1 \in H_{[\alpha]} \cap \mathbf{R}[x] \cap U_j$, $y_1 \in H_{[\alpha]} \cap \mathbf{R}[y] \cap U_j$ be points of intersection. Therefore by equation (15),

$$|\pi_A(\phi(x)) - \pi_A(\phi(x_1))| \leq \varrho \text{diam}(\mathbf{B}(\Omega_j)), \quad |\pi_A(\phi(y)) - \pi_A(\phi(y_1))| \leq \varrho \text{diam}(\mathbf{B}(\Omega_j))$$

Since $d(x_1, y_1) \leq \text{diam}(\mathbf{B}(\Omega_{j+1}))$, for $\iota = 1, 2$ we can find geodesic segments $\gamma_{x_1, \iota}, \gamma_{y_1, \iota}$ leaving x_1, y_1 respectively such that $Q = \{\gamma_{x_1, \iota}, \gamma_{y_1, \iota}\}_{\iota=1,2}$ is a 0-quadrilateral. Additionally, because $x_1, y_1 \in U_j \cap H_{[\alpha]}$, we can assume for $\iota = 1, 2$, $* = x_1, y_1$, the subsegment $\hat{\gamma}_{*, \iota} \subset \gamma_{*, \iota}$ containing $*$ and

satisfies $|\hat{\gamma}_{*,\iota}| = \frac{1}{1+\varsigma} |\gamma_{*,\iota}|$ admit geodesic approximation to its ϕ image. That is, $\phi(\hat{\gamma}_{*,\iota})$ is within $\eta|\hat{\gamma}_{*,\iota}|$ Hausdorff neighborhood of another geodesic segment.

Let $l_{*,\iota}$ be a geodesic approximation to $\phi(\hat{\gamma}_{*,\iota})$. Then angle between the direction of $l_{*,\iota}$ and that of $\hat{\gamma}_{*,\iota}$ is at most $\sin^{-1}(\hat{\eta})$. Therefore by modifying each $l_{*,\iota}$ by an amount at most $\hat{\eta}2\kappa \text{diam}(Q) \leq \hat{\eta} \text{diam}(\mathbf{B}(\Omega_{j+1}))$, we can assume $l_{*,\iota}$ all have parallel directions. Since $\varsigma \ll 1$, the four geodesic segment $l_{*,\iota}$ do constitute a quadrilateral $\tilde{Q}$ and Lemma ?? of [P] applied to $\tilde{Q}$ yields

$$|\pi_{\vec{v}} \circ \Pi_{\vec{v}}(\phi(x_1)) - \pi_{\vec{v}} \circ \Pi_{\vec{v}}(\phi(y_1))| \leq \hat{\eta} \text{diam}(\mathbf{B}(\Omega_j))$$

where $\vec{v}$ is parallel to edge directions of $\tilde{Q}$. Therefore

$$|\pi_{\vec{v}} \circ \Pi_{\vec{v}}(\phi(x)) - \pi_{\vec{v}} \circ \Pi_{\vec{v}}(\phi(y))| \leq \hat{\eta} \text{diam}(\mathbf{B}(\Omega_j)) + 2\varrho \text{diam}(\mathbf{B}(\Omega_j)) \leq 4\hat{\eta} \text{diam}(\mathbf{B}(\Omega_j))$$

since $\varrho \leq \hat{\eta}$. The claim now follows by constructing quadrilaterals whose image is approximated by quadrilaterals whose edge direction ranges over at least n many linearly independent directions. $\square$

Lemma 5.1.2. *Suppose $p \in \mathbf{R}_j[x] \cap U_j$, $q \in \mathbf{R}_{j+1}[x] \cap U_{j+1}$. Then,*

$$|\pi_A(\phi(x)) - \pi_A(\phi(y))| \leq (4|\Delta|n+2)\hat{\eta} \text{diam}(\mathbf{B}(\Omega_{j+1}))$$

Proof. Note that the projection of both $\mathbf{R}_{j+1}[x] \cap U_j$ and $\mathbf{R}_{j+1}[x] \cap U_{j+1}$ to $R_{j+1}[x]$ have relative measure up at least $1 - \varrho$. Therefore we can find $p' \in \mathbf{R}_{j+1}[x] \cap U_j$, $q' \in \mathbf{R}_{j+1}[x] \cap U_{j+1}$, and $|\pi_A(\phi(p')) - \pi_A(\phi(q'))| \leq \varrho \text{diam}(\mathbf{B}(\Omega_{j+1}))$. Now by Lemma 5.1.1,

$$|\pi_A(\phi(p)) - \pi_A(\phi(p'))| \leq (|\Delta|)n4\hat{\eta} \text{diam}(\mathbf{B}(\Omega_j))$$

and by equation (15)

$$|\pi_A(\phi(q')) - \pi_A(\phi(q))| \leq \hat{\eta} \text{diam}(\mathbf{B}(\Omega_{j+1}))$$

the claim now follows by triangle inequality. $\square$

Proof. to Theorem 5.1 We have

$$R_0[x] \subset R_1[x] \subset R_2[x] \cdots$$

and

$$R_0[y] \subset R_1[y] \subset R_2[y] \cdots$$

There exists N with $\text{diam}(\mathbf{B}(\Omega_N))$ is comparable to $d(x, y)$ (after possibly shifting the N 's grid by a bit) such that $R_N[x] = R_N[y]$. Now for $0 \leq j \leq N$, pick $x_j \in \mathbf{R}_j[x] \cap U_j$, $y_j \in \mathbf{R}_j[y] \cap U_j$. We may assume that $x_N = y_N$. By Lemma 5.1.2:

$$\begin{aligned} |\pi_A(\phi(x_0)) - \pi_A(\phi(y_0))| &\leq \sum_{j=0}^{N-1} |\pi_A(\phi(x_{j+1})) - \pi_A(\phi(x_j))| + \sum_{j=0}^{N-1} |\pi_A(\phi(y_{j+1})) - \pi_A(\phi(y_j))| \\ &\leq 2(4|\Delta|n+2) \sum_{j=0}^{N-1} \hat{\eta} \text{diam}(\mathbf{B}(\Omega_{j+1})) \\ &\leq 4(2|\Delta|n+1) \frac{\hat{\eta}}{\varsigma} \text{diam}(\mathbf{B}(\Omega_N)) \end{aligned}$$

where last inequality comes from $\text{diam}(\mathbf{B}(\Omega_{j+1})) = (1 + \varsigma)\text{diam}(\mathbf{B}(\Omega_j))$. Now since $x_0 \in R_0[x]$ $|\pi_A(\phi(x)) - \pi_A(\phi(x_0))| \leq 2\kappa d(x, x_0) \leq 2\kappa \text{diam}(\mathbf{B}(\Omega)) = M$, and similarly $|\pi_A(\phi(y)) - \pi_A(\phi(y_0))| \leq M$. The claim now follows by noting that we chose $\frac{\text{diam}(\mathbf{B}(\Omega_N))}{d(x, y)} \in [1/2, 2]$ and $\varsigma \gg \hat{\eta}$ $\square$

5.2 Consequence of weak height preservation - flats go to flats

Theorem 5.1 is the first statement we have that places no additional constraints on the points other than their natural relation in G . In this subsection we show that as a first consequence, the image of a flat is within $O(1)$ of another flat, which eventually culminating in the proof of theorem 5.2 in the next subsection.

Proposition 5.2.1. *ϕ sends each flat to within $O(1)$ of another flat.*

We now proceed to establish some necessary observations.

Lemma 5.2.1. *There is a linear map $A_0 : \mathbf{A} \rightarrow \mathbf{A}'$ and numbers $\hat{\tau} < 1$, $\hat{M} > 0$ such that for any $x, y \in G$,*

$$|(\pi_A(\phi(x)) - \pi_A(\phi(y))) - (A_0(\pi_A(x)) - A_0(\pi_A(y)))| \leq \hat{\tau}d(x, y) + \hat{M} \quad (16)$$

Proof. Let $\mathbf{B}$ be a box such that $\text{diam}(\mathbf{B})/d(x, y) \in [1/2, 2]$. By Corollary 4.0.2, there is a subset $\mathcal{P}^0 \subset \mathcal{P}(\mathbf{B})$ of relative measure at least $(1 - \sqrt{\hat{\eta}})$ that supports a standard map which is $\hat{\eta}\text{diam}(\mathbf{B})$ away from $\phi|_{\mathcal{P}^0}$. Write A_0 for the linear part of the $\mathbf{A}'$ part of the standard map. Without loss of generality we can assume that $x\mathbf{H} \cap \mathcal{P}^0 \neq \emptyset$, and $y\mathbf{H} \cap \mathcal{P}^0 \neq \emptyset$. Let $\hat{x} \in x\mathbf{H} \cap \mathcal{P}^0$, and $\hat{y} \in y\mathbf{H} \cap \mathcal{P}^0$. Then by Theorem 5.1

$$|(\pi_A(\phi(\hat{y})) - \pi_A(\phi(y))) - (A_0(\pi_A(\hat{y})) - A_0(\pi_A(y)))| \leq \tau d(\hat{y}, y) + M$$

since $\pi_A(\hat{y}) = \pi_A(y)$. By Corollary 4.0.2 we also have

$$|(\pi_A(\phi(\hat{x})) - \pi_A(\phi(\hat{y}))) - (A_0(\pi_A(\hat{x})) - A_0(\pi_A(\hat{y})))| \leq \hat{\eta}\text{diam}(\mathbf{B})$$

and by Theorem 5.1 again, we have

$$|(\pi_A(\phi(x)) - \pi_A(\phi(\hat{x}))) - (A_0(\pi_A(x)) - A_0(\pi_A(\hat{x})))| \leq \tau d(x, \hat{x}) + M$$

Summing all three equations and apply triangle inequality to the left hand side we have

$$\begin{aligned} |(\pi_A(\phi(x)) - \pi_A(\phi(y))) - (A_0(\pi_A(x)) - A_0(\pi_A(y)))| &\leq (2\tau + \eta)\text{diam}(\mathbf{B}) + 2M \\ &\leq 2(2\tau + \hat{\eta})d(x, y) + 2M \end{aligned}$$

since $\hat{\eta}$ and τ depends on our initial ϵ, δ , and approach zero as those initial input approach zero, we can assume that $2(2\tau + \hat{\eta}) < 1$, and we set $\hat{\tau} = 2(2\tau + \hat{\eta})$, $\hat{M} = 2M$. $\square$

Corollary 5.2.1. *There is a number M_0 such that if $p, q \in G$ are two points on the same flat and $\pi_A(\phi(p)) = \pi_A(\phi(q))$, then $d(x, y) \leq M_0$.*

Proof. In equation (16) substitute x, y by p, q and note that since p, q lies on the same flat, $|\pi_A(p) - \pi_A(q)| = d(p, q)$. So we have

$$(1 - \hat{\tau})d(p, q) \leq \hat{M}$$

to which the result follows. Alternatively, this can be obtained from Theorem 5.1 applied to the inverse map ϕ^{-1} and $\phi(p)$, $\phi(q)$, for then equation (14) becomes

$$\begin{aligned} |\pi_A \circ \phi^{-1}(\phi(p)) - \pi_A \circ \phi^{-1}(\phi(q))| &\leq \tau d(\phi(p), \phi(q)) + M \leq \tau \kappa d(p, q) + M + C \\ |\pi_A(p) - \pi_A(q)| = d(p, q) &\leq \tau \kappa d(p, q) + M + C \\ d(p, q) &\leq \frac{M + C}{1 - \tau \kappa} \end{aligned}$$

□

A subset L of $\mathbf{A}' \simeq \mathbb{R}^n$ is called a 'grid' if it is the image of an injective homomorphism $\psi : \mathbb{Z}^n \rightarrow \mathbf{A}'$. A line in L refers to a subset of the form: $\{\psi(\mathbf{c} + t\mathbf{u}) : t \in \mathbb{Z}\}$ for some $\mathbf{c}, \mathbf{u} \in \mathbb{Z}^n$, and each coordinate of $\mathbf{u}$ is either $+1, -1$ or 0 . A grid is said to be good if none of its lines are parallel to root kernels.

Lemma 5.2.2. *Let $\{x_i\} \subset G'$ be a sequence of points with the following properties:*

- (i) $\pi_A(x_j) \neq \pi_A(x_i)$ if $i \neq j$.
- (ii) $\{\pi_A(x_i)\} \subset \mathbf{A}'$ is a good grid.
- (iii) for any subsequence $\{x_{i_j}\}$ such that $\{\pi_A(x_{i_j})\}$ is a line, $\{x_{i_j}\}$ is within $O(1)$ of a (bi-infinite) geodesic.

Then $\{x_i\}$ is within $O(1)$ of a flat.

Proof. Write $\{\pi_A(x_i)\} = L$. Let $\{x_{1_j}\}$, and $\{x_{2_j}\}$ be two subsequences such that their π_A images are two parallel lines, and let l_1, l_2 be two geodesics within $O(1)$ of $\{x_{1_j}\}$ and $\{x_{2_j}\}$ respectively. We note to every x_{1_j} , there is a x_{2_j} such that $\pi_A(x_{2_j})$ is closest to $\pi_A(x_{1_j})$ amongst $\{\pi_A(x_{2_j})\}$, and furthermore, there is a line in L containing $\pi_A(x_{1_j})$ and $\pi_A(x_{2_j})$, and by assumption, this means that there is a geodesic $\hat{l}_j$ (whose direction is the same for all j) within $O(1)$ of x_{1_j} and x_{2_j} . Now if l_1 and l_2 are not in the same flat, then for any root Ξ , $\pi_\Xi(l_1)$ and $\pi_\Xi(l_2)$ fork out with respect to the orientation we previously fixed on l_1 and l_2 . When x_{1_j}, x_{2_j} are far away from the fork point, this causes a contradiction because the existence of $\hat{l}_j$ mean that $\pi_\Xi(x_{1_j})$ and $\pi_\Xi(x_{2_j})$ can be connected by a vertical geodesic in V_Ξ , but they lie far away from the forking point of two geodesics. Therefore if $\{x_{i_j}\}$ is a subsequence for which their π_A image is a affine 2-subspace, then $\{x_{i_j}\}$ lie within $O(1)$ of an affine 2-subspace in a flat.

Now suppose whenever $\{x_{i_j}\}$ is a subset whose π_A image is an affine I -subspace, $\{x_{i_j}\}$ is within $O(1)$ of a flat. Let $\{x_{1_j}\}, \{x_{2_j}\}$ be two subsets such that $\{\pi_A(x_{1_j})\}, \{\pi_A(x_{2_j})\}$ are two parallel I -hyperplane, and h_1, h_2 be two affine I subspace within $O(1)$ of $\{x_{1_j}\}$ and $\{x_{2_j}\}$ respectively.

Then we know for every x_{1_j} , there is a x_{2_j} such that there is a line in L containing $\pi_A(x_{1_j})$ and $\pi_A(x_{2_j})$, so by assumption, x_{1_j}, x_{2_j} are within $O(1)$ of a (straight) geodesic $\hat{l}_j$. Furthermore we can assume WLOG that the direction of $\hat{l}_j$ are the same for all j . Therefore if h_1 and h_2 lie on two different flats, then for some root Ξ , $\pi_\Xi(h_1)$ and $\pi_\Xi(h_2)$ are two vertical geodesic that fork apart. Therefore for x_{1_j}, x_{2_j} such that $\pi_\Xi(x_{1_j}), \pi_\Xi(x_{2_j})$ lying very far from the fork point where $\pi_\Xi(h_1)$ and $\pi_\Xi(h_2)$ diverge from each other, this is a contradiction to the existence of $\hat{l}_j$ within $O(1)$ of x_{1_j} and x_{2_j} , for the latter would imply that $\pi_\Xi(x_{1_j})$ and $\pi_\Xi(x_{2_j})$ are within $O(1)$ of a vertical geodesic in V_Ξ . □

Proof. to Proposition 5.2.1

Fix a flat $\mathcal{F} \simeq \mathbf{A}$. The composition $\pi_A \circ \phi|_{\mathcal{F}}$ is homotopic to the identity map when we consider $\mathcal{F}$ as $\mathbf{A}$, so $\pi_A \circ \phi|_{\mathcal{F}}$ preserves homologies. By equation (16), it takes balls to neighborhood of balls, so it preserves homologies relative to the complement of a ball, therefore $\pi_A \circ \phi|_{\mathcal{F}}$ is onto.

Let L be a good grid in $\mathbf{A}'$ under a group isomorphism ψ . We can further make sure that for each basis $e_i \in \mathbb{Z}^n$, $|\psi(e_i) - \psi(\vec{0})| = 4M_0$, where M_0 is as in Corollary 5.2.1. The same corollary also implies that for every point $\mathbf{b} \in L$, the subset $s_{\mathbf{b}} = \{x \in \phi(\mathcal{F}) : \pi_A(x) = \mathbf{b}\}$ of G' is contained in a ball of radius at most M_0 . Therefore for distinct points $\mathbf{b}, \mathbf{d} \in L$, $s_{\mathbf{b}} \cap s_{\mathbf{d}} = \emptyset$.

Let $\{x_j\} \subset \phi(\mathcal{F})$ be a subset such that $x_j \in s_{\mathbf{b}}$ for some $\mathbf{b} \in L$. We choose x_j 's so that if $j \neq i$, $x_j \in s_{\mathbf{b}}$, $x_i \in s_{\mathbf{c}}$ for $\mathbf{b} \neq \mathbf{c}$. Now let $\{x_{i_j}\}$ be a subsequence whose π_A images is a line in L , and $\{y_{i_j}\} \subset \mathcal{F}$ be their ϕ -preimages in $\mathcal{F}$. Then by equation (16), $d(y_{i_j}, y_{i_{j+1}}) \leq \hat{C}(L)$, for all j , for some constant $\hat{C}$ depending on L . This means that that $d(x_{i_j}, x_{i_{j+1}}) \leq 2\kappa\hat{C}$ for all j . Apply (i) of Lemma ?? to $\{x_{i_j}\}$ shows that the sequence is within $O(\hat{C})$ of a (bi-infinite) geodesic. The claim now follows by applying Lemma 5.2.2 to $\{x_j\}$. $\square$

5.3 Consequences of flats go to flats

The primarily reason for introducing root boundaries $\partial_{[\alpha]}$ for each root class $[\alpha]$ and the (psudo) distance function is that a quasi-isometry of G induces quasi-similarities among $(\partial_{[\alpha]}, D_{[\alpha]})$'s.

Recall that a map $F : X \rightarrow Y$ between metric spaces is called a (N, K) -quasi-similarities if

$$N/Kd(x, y) \leq d(F(x), F(y)) \leq NKd(x, y)$$

When $K = 1$, the map is called a *similarity*. We write the group of quasi-similarities of a space by $QSim(X)$.

Proposition 5.3.1. *If $\psi : G = \mathbf{H} \rtimes_{\varphi} \mathbf{A} \rightarrow G' = \mathbf{H}' \rtimes_{\varphi'} \mathbf{A}'$ is a quasi-isometry such that the image of left translate of $\mathbf{A}$ is within $O(1)$ neighborhood of a left translates of $\mathbf{A}'$. Then*

- (i) ψ sends foliations by root kernels of G to the foliations by root kernels of G' .
- (ii) The number of root classes of G and G' are the same.
- (iii) ψ sends foliations by root class horocycles of G to foliations by root class horocycles of G' .
- (iv) ψ is within $O(1)$ of a product map $f \times g$, where $f : \mathbf{A} \rightarrow \mathbf{A}'$ is affine whose linear part is some scalar multiple of a finite order element of $O(n)$; while $g = (g_1, g_2, \dots, g_{s_0})$, each g_i is a bilipschitz map from $V_{[\alpha]}$ to $V_{f_*[\alpha]}$, where f_* is the bijection from root classes of G to root classes of G' as induced by the linear part of f .
- (v) For each root class $[\Xi]$, if we list the roots $\xi_1 < \xi_2 < \xi_3 \dots \xi_l$, then f induces a map on the roots such that $f_*\xi_1 < f_*\xi_2 \dots f_*\xi_l$, where $\{f_*\xi\} = f_*[\Xi]$. Furthermore, with respect to this order of roots, $g|_{V_{[\Xi]}} : V_{[\Xi]} \rightarrow V_{f_*[\Xi]}$ has the form

$$(x_1, x_2, \dots, x_l) \mapsto (g_1(x_1, x_2, \dots, x_l), \dots, g_{l-1}(x_{l-1}, x_l), g_l(x_l))$$

such that for each i , any $x_j \in V_{\xi_j}$ for $j > i$, the map $\bullet \mapsto g_i(\bullet, x_{i+1}, \dots, x_l)$ is a $(e^{f_*\xi_i(t_0)}, 2\kappa e^c)$ quasi-similarity, where t_0 is the constant part of f , and c is a constant depending on ψ .

(vi) ψ induces an quasi-similarity between root boundaries of G to root boundaries of G' .

Proof. Since two flats are within a finite Hausdorff distance of each other if and only if they are the same flat, our assumption means that a flat $\hat{\mathcal{F}}$ within $O(1)$ Hausdorff neighborhood of $\psi(\mathcal{F})$ for $\mathcal{F}$ a flat, is unique.

If two flats $\mathcal{F}_1, \mathcal{F}_2$ have no-empty intersection (i.e. come together) then $\hat{\mathcal{F}}_1 \cap \hat{\mathcal{F}}_2 \neq \emptyset$, and $d_H(\phi(\partial(\mathcal{F}_1 \cap \mathcal{F}_2)), \partial(\hat{\mathcal{F}}_1 \cap \hat{\mathcal{F}}_2)) \leq M_0$.

The boundary of the set where two flats come together viewed as a subset of $\mathbf{A}$ is a finite union of hyperplanes parallel root kernels. If for two flats $\mathcal{F}, \mathcal{F}'$, $\partial(\mathcal{F} \cap \mathcal{F}')$ is a union of hyperplanes parallel to at least two roots kernels, then there are flats $\mathcal{F}_1, \mathcal{F}_2, \dots, \mathcal{F}_k$ such that $\mathcal{F} \cap \mathcal{F}' = \bigcap_{i=1}^k \mathcal{F} \cap \mathcal{F}_i$.

Therefore if $\partial(\mathcal{F}_1 \cap \mathcal{F}_2)$ is a hyperplane parallel to just one root kernel, then so is $\partial(\hat{\mathcal{F}}_1 \cap \hat{\mathcal{F}}_2)$.

Fix a flat $\mathcal{F}$. Since every hyperplane in $\mathcal{F}$ parallel to a root kernel is the boundary of $\mathcal{F} \cap \mathcal{F}'$ for some other flat $\mathcal{F}'$, it follows that if $h_1, h_2 \subset \mathcal{F}$ are two hyperplanes parallel to a common root kernel, then there exists hyperplanes $\hat{h}_1, \hat{h}_2 \subset \hat{\mathcal{F}}$ parallel to a common root kernel such that $d_H(\phi(h_i), \hat{h}_i) \leq A$ for $i = 1, 2$. In other words, $\phi|_{\mathcal{F}} : \mathcal{F} \rightarrow \hat{\mathcal{F}}$ sends hyperplanes parallel to the kernel of a root, to hyperplanes parallel to the kernel of some other root. As ψ is coarsely onto, it follows that the number of root kernels of G and G' are the same.

For the remaining claims, we consider two cases separately.

Case I - G has rank(G) many root kernel

Since G is unimodular, the sum of all its roots is zero. But with only $\text{rank}(G)$ many root kernels, this means that there is a basis $\{\alpha_i\}$ of $\mathbf{A}^*$ such that every root of G is of the form $c\alpha_i$ where $c \in \mathbb{R} - \{0\}$. By abuse of notation, we will write $V_{\alpha_i}^+$ to be the direct sum of root spaces where the root is positive multiples of α_i , even though α_i itself might not be a root, and similarly, $V_{\alpha_i}^-$ denote for direct sum of root spaces where the root is negative multiples of α_i . Let $\{\beta_i\} \subset \mathbf{A}'^*$ be the analogous linear functionals for G' . In this way, $\mathbf{H} = \bigoplus_i V_{\alpha_i}^+ \oplus V_{\alpha_i}^-$, and similarly $\mathbf{H}' = \bigoplus_i V_{\beta_i}^+ \oplus V_{\beta_i}^-$.

First we show that left translates of $\mathbf{H}$ are taken to $O(1)$ Hausdorff neighborhood of left translates of $\mathbf{H}'$. Take $p, q \in V_{\alpha_1}^*$, $\star \in \{+, -\}$. We construct a quadrilateral Q (see Definition ?? in [P]) with p, q as two of its vertices using left translates of $\bigcap_{j \neq 1} \ker(\alpha_j)$. Since flats are taken to $O(1)$ neighborhoods of flats, $\psi(Q)$ is within $O(1)$ neighborhood of a quadrilateral $\hat{Q}$ whose edges are left translates of $\bigcap_{j \neq l} \ker(\beta_j)$ for some l , and therefore $\psi(p), \psi(q)$ are within $O(1)$ neighborhood of a left translate of $V_{\beta_l}^*$, $\star \in \{+, -\}$.

More generally, for any two points $x, y \in \mathbf{H}$, we can find a sequence of points $p_i \in \mathbf{H}$, such that $p_0 = x$, $p_s = y$, and for all i , p_i, p_{i+1} differ only in a $V_{\alpha_i}^*$, $\star \in \{+, -\}$ coordinate. The previous argument then implies $\psi(x)$ and $\psi(y)$ are within $O(1)$ of a left translate of $\mathbf{H}'$, because the same is true of $\psi(p_i)$ and $\psi(p_{i+1})$ for all i .

Next we show that the restriction of ψ to different flats agree. If $p, q \in \mathbf{H}$ are points such that for each α_i , these two points differ in only one of $V_{\alpha_i}^+, V_{\alpha_i}^-$ coordinate (but not both), then the intersection of flats based on them, $p\mathcal{F}, q\mathcal{F}$, intersect at an unbounded set whose boundary is a union of hyperplane parallel to kernels of α_i 's that appear in the coordinate difference between p and q . Furthermore, we can find a third point $x \in \mathbf{H}$ that differ in one of $V_{\alpha_i}^+, V_{\alpha_i}^-$ coordinate with each of p and q so that $p\mathcal{F} \cap q\mathcal{F} \subset p\mathcal{F} \cap x\mathcal{F}$, and $p\mathcal{F} \cap q\mathcal{F} \subset q\mathcal{F} \cap x\mathcal{F}$. In this way, we see that $\phi|_{p\mathcal{F}}$ is the same as $\phi|_{q\mathcal{F}}$.

More generally, for any two points $x, y \in \mathbf{H}$, we can find a finite number of points $p_i \in \mathbf{H}$ such that $p_0 = x$, $p_l = y$, such that for all i , p_i differ from p_{i+1} in one of $V_{\alpha_j}^+$, $V_{\alpha_j}^-$ coordinate (but not both). The previous argument then implies $\phi|_{x\mathcal{F}}$ is the same as $\phi|_{x\mathcal{F}}$ (as $\phi|_{p_i\mathcal{F}}$ is the same as $\phi|_{p_{i+1}\mathcal{F}}$).

Denote the common value of the restriction of ψ to any flat as f . Then we know that f induces a bijection between $\{\alpha_i\}$ and $\{\beta_i\}$ which are numbered such that for all i , ψ sends left translates of kernel of α_i to left translates of kernel of β_i . As left translate of $\ker(\alpha_i)$ (resp. $\ker(\beta_i)$) can be identified with a rank 1 space $G_{\alpha_i} = \mathbf{H} \rtimes \mathbb{R}(\vec{v}_{\alpha_i})$ (resp. $G'_{\beta_i} = \mathbf{H}' \rtimes \mathbb{R}(\vec{v}_{\beta_i})$), ψ induces a quasi-isometry from G_{α_i} to G'_{β_i} that sends left translates of $\mathbf{H}$ to $O(1)$ neighborhood of left translates of G'_{β_i} . By Proposition 5.8 of [FM], we conclude that the f -induced map from $\mathbf{A}/\ker(\alpha_i)$ to $\mathbf{A}'/\ker(\beta_i)$ is bounded distance from an affine map.

Repeat the above argument to all other α_i 's shows that f is $O(1)$ away from being affine that respects root kernels of G and G' .

Case II - G has at least $\text{rank}(\mathbf{G})+1$ many root kernels. In this case, the restriction to a flat, $\phi|_{\mathcal{F}}$ preserves at least $n+1$ many parallel families of hyperplanes, thus forcing $\phi|_{\mathcal{F}}$ to be affine (when $\mathcal{F}$ and $\hat{\mathcal{F}}$ are viewed as $\mathbf{A}$ and $\mathbf{A}'$ respectively.) that respects root kernels.

So by identifying $\mathcal{F}$ and $\hat{\mathcal{F}}$ with $\mathbb{R}^n$, $\phi|_{\mathcal{F}}$ can be written as $\mathbf{t} \mapsto M_{\mathcal{F}}(\mathbf{t}) + \mathbf{t}_0^{\mathcal{F}}$, where $M_{\mathcal{F}} \in GL(n)$ is a linear map preserving root kernels. This implies (i), that images of (straight) geodesics are within $O(1)$ Hausdorff neighborhood of straight geodesics. At this stage, both the affine map $M_{\mathcal{F}}$ and the translation $\mathbf{t}_0^{\mathcal{F}}$ depend on the flat $\mathcal{F}$.

Fix a root equivalence class $[\Xi]$. Let $\vec{v} \in \mathbb{R}^n$ be a direction such that $\alpha(\vec{v}) \neq 0$ for all roots α . Then for any two points x, y in the same left coset of $V_{[\Xi]}$, there is a quadrilateral Q with x, y as two of its vertices, and whose edges are all in direction $\vec{v}$. WLOG we can assume $\Xi(\vec{v}) < 0$, so x, y are in the same left coset of $W_{\vec{v}}^-$. Since ψ takes geodesics to $O(1)$ neighborhood of geodesics, it follows that $\psi(Q)$ is within $O(1)$ of another (0) quadrilateral. Write $\mathcal{F}_x = x\mathbf{A}$, $\mathcal{F}_y = y\mathbf{A}$ for the flats based at x and y respectively. Then $\psi(Q)$ close to a quadrilateral means that $M_{\mathcal{F}_x}(\vec{v}) = M_{\mathcal{F}_y}(\vec{v})$, and the dot product between $\vec{v}$ and $\mathbf{t}_0^{\mathcal{F}_x}$, as well as $\mathbf{t}_0^{\mathcal{F}_y}$ are the same. More over, writing $\vec{u} = M_{\mathcal{F}_x}(\vec{v})$, by Lemma ?? in [P], $\psi(x)$, $\psi(y)$ are within $O(1)$ of a left coset of $W_{\vec{u}}^-$.

We can find at least n $\vec{v}_i$'s, such that $\cap_i W_{\vec{v}_i}^{\Xi(\vec{v}_i)/|\Xi(\vec{v}_i)|} = V_{[\Xi]}$ and none of them lies in a root kernel. Repeat the same argument as before yields that $M_{\mathcal{F}_x} = M_{\mathcal{F}_y} = M$, $\mathbf{t}_0^{\mathcal{F}_x} = \mathbf{t}_0^{\mathcal{F}_y}$. That is, the restriction of ψ to the flats based at x, y are within $O(1)$ of each other. WLOG, we can assume that we have so many $\vec{v}_i$'s, such that $\cap_i W_{\vec{v}_i}^{\Xi(\vec{v}_i)/|\Xi(\vec{v}_i)|} = V_{[\beta]}$ for some root equivalence class $[\beta]$, which shows that $\psi(x)$, $\psi(y)$ are within $O(1)$ of a left translate of $V_{[\beta]}$ for some root class $[\beta]$.

Since any two points on the same left translate of $\mathbb{R}^m$ (or $\mathbf{H}$) can be 'connected' by finitely many points, for which successive pairs of points lie on a left translate of $V_{[\alpha]}$ for some root class $[\alpha]$, it follows that ψ sends a left translate of $\mathbf{H}$ to within $O(1)$ neighborhood of another left translate of $\mathbf{H}$. Since the ψ restricted to flats based at two points that lie on a common horocycle agree, we also have now that the restriction of ψ to any two flats agree. That is, the restriction of ψ to any flat does not depend on the base point of the flat.

So now we know that regardless of the number of root kernels, ψ splits into $f \times g$, where $f : \mathbf{A} \rightarrow \mathbf{A}'$ affine and respects root kernels, while $g : \mathbf{H} \rightarrow \mathbf{H}'$ takes root class horocycles to root class horocycles. Furthermore, the permutation on root classes induced by f and g agree.

We now proceed to show that the f actually induces a bijection between roots of G and G' (not just root classes).

Since we know now the ψ restricted to any flat is the map $\mathbf{t} \mapsto M(\mathbf{t}) + \mathbf{t}_0$. This means that (straight) geodesics are taken to straight geodesics, and we can compare the rate of divergence between two geodesics in the same direction but at based at different points of $\mathbf{H}$.

Specifically, take Ξ a root, and let p, q be two points on a common Ξ horocycle. Pick some $\vec{v} \in \mathbf{A}$, and let $l_x = x(t\vec{v})$, $l_y = y(t\vec{v})$ be two geodesic rays in direction $\vec{v}$ leaving x, y respectively. Then $\psi(l_x)$, $\psi(l_y)$ are within $O(1)$ of $l_{x'} = x'(tM(\vec{v}))$ and $l_{y'} = y'(tM(\vec{v}))$ respectively, where x', y' are within $O(1)$ of a left translate of $V_{[\beta]}$.

The rate of divergence between l_x and l_y is $e^{t\Xi(\vec{v})}$; the divergence rate between $l_{x'}$ and $l_{y'}$ is $e^{t\xi(M(\vec{v}))}$ for some $\xi \in [\beta]$. Since the two rates are QI to each other, there must be some $\tilde{\beta} \in [\beta]$ such that $\Xi(\vec{v}) = \tilde{\beta}(M(\vec{v}))$. But $\vec{v}$ is arbitrary, so $\Xi = \tilde{\beta} \circ M$, x', y' are in the same left translate of $\oplus_{\xi \in [\beta]: \xi < \tilde{\beta}} V_\xi$.

This means that f induces a bijection σ_f between roots of G and G' . So that for each root α , ψ sends left translates of $\oplus_{\xi \in [\alpha]: \xi \leq \alpha} V_\xi$, to $O(1)$ Hausdorff neighborhoods of left translates of direct sum of root spaces where the roots are in the same root class as $\sigma_f(\alpha)$ but less than $\sigma_f(\alpha)$.

For $x, y \in V_\Xi$, write $g_\Xi(x), g_\Xi(y) \in V_{\sigma(\Xi)}$ for the $V_{\sigma(\Xi)}$, and for each $\xi < \sigma(\Xi)$, write $\tilde{g}_\xi(x), \tilde{g}_\xi(y) \in V_\xi$ for the V_ξ component of $g(x)$ and $g(y)$, so that $g(x) = g_\Xi(x) + \sum_{\xi < \sigma(\Xi)} \tilde{g}_\xi(x)$, $g(y) = g_\Xi(y) + \sum_{\xi < \sigma(\Xi)} \tilde{g}_\xi(y)$. Pick a $\mathbf{t} \in \mathbf{A}$, then the distance between $(\mathbf{t}, x)$ and $(\mathbf{t}, y)$ with respect to path metric in $\mathbf{tH}$ is $e^{-\Xi(\mathbf{t})}|x - y|$. The ψ images of $(\mathbf{t}, x)$, $(\mathbf{t}, y)$ is c away from $(M\mathbf{t} + \mathbf{t}_0, g(x))$ and $(M\mathbf{t} + \mathbf{t}_0, g(y))$, so we have the following inequality:

$$\begin{aligned} \frac{1}{2\kappa} e^{-c} e^{-\Xi(\mathbf{t})} |x - y| &\leq \left(e^{-\sigma(\Xi)(M\mathbf{t} + \mathbf{t}_0)} P_\Xi(\sigma(\Xi)(M\mathbf{t} + \mathbf{t}_0)) |g_\Xi(x) - g_\Xi(y)| \right) \\ &\quad + \left(\sum_{\xi < \sigma(\Xi)} e^{-\xi(M\mathbf{t} + \mathbf{t}_0)} P_\xi(\xi(M\mathbf{t} + \mathbf{t}_0)) |\tilde{g}_\xi(x) - \tilde{g}_\xi(y)| \right) \\ &\leq 2\kappa e^c e^{-\Xi(\mathbf{t})} |x - y| \end{aligned}$$

Since $\sigma(\Xi) \circ M = \Xi$, dividing by ΞP_Ξ on both sides and let $\Xi(\mathbf{t}) \rightarrow \infty$, and noting that for any polynomial Q , $\lim_{t \rightarrow \infty} Q(t + t_0)/Q(t)$ is bounded above by a number independent of t_0 , so we end up with

$$\frac{1}{2\kappa} e^{-c} e^{\sigma(\Xi)(\mathbf{t}_0)} |x - y| \leq |g_\Xi(x) - g_\Xi(y)| \leq 2\kappa e^c e^{\sigma(\Xi)(\mathbf{t}_0)} |x - y|$$

so the restriction of $g|_{V_\Xi} : V_\Xi \rightarrow V_{\sigma(\Xi)}$ is bilip with bilip constants $2\kappa e^c e^{\sigma(\Xi)(\mathbf{t}_0)}$, $1/2\kappa e^{-c} e^{\sigma(\Xi)(\mathbf{t}_0)}$.

To summarize, ψ is $O(1)$ from a map of the form $(\mathbf{x}, \mathbf{t}) \mapsto (g(\mathbf{x}), m A_f(\mathbf{t}) + \mathbf{t}_0)$, where $m > 0$, $A_f : \mathbf{A} \rightarrow \mathbf{A}'$ is a finite order element in $O(n)$ that preserves foliations by root kernels, while $g : \mathbf{H} \rightarrow \mathbf{H}'$ sends root horocycles to root horocycles and furthermore respects the graded foliations in each root horocycle.

Let σ be the permutation on root classes induced by M . Then as ψ sends negative $[\alpha]$ half planes to bounded neighborhood of negative $\sigma([\alpha])$ half planes, ψ induces a map from $\partial_{[\alpha]}^-$ to $\partial_{\sigma([\alpha])}^-$ for every root class $[\alpha]$ of G .

Furthermore, the map $\hat{q} : \mathbb{R} \sim \mathbf{A}/\ker(\alpha_0) \rightarrow \mathbf{A}/\ker(\sigma(\alpha)_0) \sim \mathbb{R}$ is bounded distance from an affine with linear term as m and constant term as $\sigma(\alpha)_0(\mathbf{t}_0)$, we now show the induced map on lower root boundaries is a quasi-similarity.

Take $p, q \in \partial_{[\alpha]} \sim V_{[\alpha]}$. Then, definition of $D_{[\alpha]}(p, q)$ says that

$$D_{[\alpha]}(p, q) = e^{mt_{p,q}}$$

under ϕ , we have

$$D_{\sigma([\alpha])}(g(p), g(q)) = e^{\hat{q}(t_{p,q})}$$

therefore

$$\begin{aligned} mt_{p,q} + \sigma(\alpha)_0(\mathbf{t}_0) - c &\leq \hat{q}(t_{p,q}) \leq mt_{p,q} + \sigma(\alpha)_0(\mathbf{t}_0) + c \\ e^{mt_{p,q}} \frac{e^{\sigma(\alpha)_0(\mathbf{t}_0)}}{e^c} &\leq e^{\hat{q}(t_{p,q})} \leq e^{mt_{p,q}} e^{\sigma(\alpha)_0(\mathbf{t}_0)} e^c \\ \frac{1}{e^c} \left(e^{\sigma(\alpha)_0(\mathbf{t}_0)} D_{[\alpha]}(p, q) \right) &\leq D_{\sigma([\alpha])}(g(p), g(q)) \leq e^c \left(e^{\sigma(\alpha)_0(\mathbf{t}_0)} D_{[\alpha]}(p, q) \right) \end{aligned}$$

□

Theorem 5.2. *Let G, G' be a non-degenerate, unimodular, split abelian-by-abelian solvable Lie group, and $\phi : G \rightarrow G'$ a (κ, C) quasi-isometry. Then ϕ is bounded distance from a composition of a left translation followed by a map of the form $(\mathbf{x}, \mathbf{t}) \rightarrow (g(\mathbf{x}), f(\mathbf{t}))$, where f is affine whose linear part is a positive of a finite order element $A_f \in O(n)$ (n is the rank of G) that preserves foliations by root kernels, while $g = (g_1, g_2, \dots, g_d)$, g_i is a bilip map from $V_{[\alpha_i]}$ to $V_{[\alpha_i]}$ with bilip constants depending only on κ, C .*

Proof. Apply Proposition 5.3.1 in light of Proposition 5.2.1. □

Corollary 5.3.1. *Let $G = \mathbf{H} \rtimes_{\varphi} \mathbf{A}$ be a non-degenerate, unimodular, split abelian-by-abelian group such that $\varphi(\mathbf{A})$ is diagonalizable, while $G' = \mathbf{H}' \rtimes_{\varphi'} \mathbf{A}'$ is another non-degenerate, unimodular, split abelian-by-abelian group such that $\varphi'(\mathbf{A}')$ is not diagonalizable. Then G and G' are not quasi-isometric.*

Proof. If there were, then Theorem 5.2 implies that geodesics are taken to geodesics and the induced height function on the geodesics are affine, which means that we can compare rates of divergence between two geodesics in the same direction. In G' , we would detect exponential polynomial growth while in G , only exponential growth can be detected, and those two growth types are not q.i. to each other. □

When the homomorphism appeared in the semidirect expression of a non-degenerate, unimodular, abelian-by-abelian solvable Lie group is diagonalizable, uniform subgroups of quasi-similarities of its root boundaries are analyzed in [D], and in this case we are able to say something about an arbitrary finitely generated group in its quasi-isometric class.

Corollary 5.3.2. *Let $G = \mathbf{H} \rtimes_{\psi} \mathbf{A}$ be a non-degenerate, unimodular, split abelian-by-abelian solvable Lie group where ψ is diagonalizable. If Γ is a finitely generated group quasi-isometric to G , then Γ is virtually polycyclic.*

Proof. Let $\varphi : \Gamma \rightarrow G$ be a (κ, C) quasi-isometry. For each $\gamma \in \Gamma$, write L_γ for the left translation of γ , and $\tilde{L}_\gamma = \varphi \circ L_\gamma \circ \varphi^{-1}$. Then $\tilde{\Gamma} = \{\tilde{L}_\gamma\}_{\gamma \in \Gamma}$ constitute a uniform subgroup of $QI(G)$, all with the same q.i. constants (they are all (κ, C) quasi-isometries).

By Theorem 5.2, each $\tilde{L}_\gamma$ induces a permutation on root classes. Therefore the map from $\tilde{\Gamma}$ into the permutations on root classes of G is a well-defined homomorphism, whose kernel, $\tilde{\Gamma}_0$ is finite index in $\tilde{\Gamma}$.

Since $\tilde{\Gamma}$ is a uniform subgroup of $QI(G)$, by Proposition 5.3.1 $\tilde{\Gamma}_0$ is a uniform subgroup of $\prod_{[\alpha]} QSim(\partial_{[\alpha]}^-)$. Applying Theorem 2 in [D] to the image of $\tilde{\Gamma}_0$ in each $QSim(\partial_{[\alpha]}^-)$ factor, we can conjugate $\tilde{\Gamma}_0$ into $\prod_{[\alpha]} ASim(\partial_{[\alpha]}^-)$, the group of almost similarities. Denote the image of $\tilde{\Gamma}_0$ in $\prod_{[\alpha]} ASim(\partial_{[\alpha]}^-)$ by $\hat{\Gamma}_0$. Note that $\hat{\Gamma}_0$ and $\tilde{\Gamma}_0$ are isomorphic.

For each $\tilde{L}_\gamma \in \tilde{\Gamma}_0$, write $g_{[\Xi], \gamma}$ for the almost similarity on $(\partial_{[\Xi]}^-, D_{[\Xi]})$ and $t_{[\Xi], \gamma}$ the corresponding similarity constant, as induced by the image of $\tilde{L}_\gamma$ in $\hat{\Gamma}_0$.

Claim: For each $\tilde{L}_\gamma \in \tilde{\Gamma}_0$, there is a $\mathbf{s}_\gamma \in \mathbf{A}$ such that $t_{[\Xi], \gamma} = e^{\Xi_0(\mathbf{s}_\gamma)}$ for each root class $[\Xi]$.

We know that for each $\tilde{L}_\gamma$, there is a $\mathbf{t}_{0, \gamma} \in \mathbf{A}$ such that $\tilde{L}_\gamma$ induces $(e^{\Xi_0(\mathbf{t}_{0, \gamma})}, e^c)$ quasi-similarity $\tilde{g}_{[\Xi], \gamma}$ on $(\partial_{[\Xi]}^-, D_{[\Xi]})$. Theorem 2 of [D] says that we can find $F_{[\Xi]} \in Bilip(\partial_{[\Xi]}^-, D_{[\Xi]})$ with bilip constant K' such that for every root class $[\Xi]$ and every $\tilde{L}_\gamma \in \tilde{\Gamma}_0$,

$$F_{[\Xi]} \tilde{g}_{[\Xi], \gamma} F_{[\Xi]}^{-1} = g_{[\Xi], \gamma}$$

Therefore $t_{[\Xi], \gamma}$, the similarity constant of $g_{[\Xi], \gamma}$ satisfies

$$t_{[\Xi], \gamma} \in [e^{\Xi_0(\mathbf{t}_{0, \gamma})} \frac{1}{e^c K'^2}, e^{\Xi_0(\mathbf{t}_{0, \gamma})} e^c K'^2], \text{ for all root classes } [\Xi] \quad (17)$$

Since the sum of roots is zero and

$$0 = \sum_{\alpha \text{ roots}} \alpha = \sum_{[\Xi]} \Xi_0 \frac{l_{[\Xi]}}{\Xi_0}$$

it follows that

$$\prod_{[\Xi]} \left(e^{\Xi_0(\mathbf{t}_{0, \gamma})} \right)^{\frac{l_{[\Xi]}}{\Xi_0}} = 1$$

therefore

$$\prod_{[\Xi]} (t_{[\Xi], \gamma})^{\frac{l_{[\Xi]}}{\Xi_0}} = 1 \quad (18)$$

(because the left hand side lies in an interval those end points are constants independent of γ , so if the left hand side was not 1, then the product for sufficiently high powers of γ escape the interval)

On the other hand, for a generic linear functional ℓ , we know that

$$0 = \sum_{[\Xi]} l_{[\Xi]}(\vec{v}_\ell) = \sum_{[\Xi]} \Xi_0(\vec{v}_\ell) \frac{l_{[\Xi]}}{\Xi_0} = \sum_{[\Xi]} \Xi_0(\mathbf{t}_{0, \gamma}) \frac{\Xi_0(\vec{v}_\ell)}{\Xi_0(\mathbf{t}_{0, \gamma})} \frac{l_{[\Xi]}}{\Xi_0}$$

By equation (17) this means that

$$\prod_{[\Xi]} (t_{[\Xi], \gamma})^{\frac{\Xi_0(\bar{v}_\ell)}{\Xi_0(\mathbf{t}_{0, \gamma})} \frac{t_{[\Xi]}}{\Xi_0}} = 1 \quad (19)$$

By letting ℓ ranging over a subset of positive measure, equation (19) and (18) means that the $t_{[\Xi], \gamma}$ must be of the form $e^{\Xi_0(\mathbf{s}_\gamma)}$ for some $\mathbf{s}_\gamma \in \mathbf{A}^2$.

This means that for each $\tilde{L}_\gamma \in \tilde{\Gamma}_0$, $\{g_{[\Xi], \gamma}\}_{[\Xi]}$ determines an element $\psi_\gamma \in QI(G)$ of the form

$$\psi_\gamma((\mathbf{x}_{[\Xi]})_{[\Xi]}, \mathbf{t}) = ((g_{[\Xi], \gamma}(\mathbf{x}_{[\Xi]}))_{[\Xi]}, \mathbf{t} + \mathbf{s}_\gamma)$$

and we can define a homomorphism $h : \hat{\Gamma}_0 \rightarrow \mathbf{A}$ as $\gamma \mapsto \mathbf{s}_\gamma$.

The kernel of h consist of elements with no translations, so they leave the subgroup $\mathbf{H}$ invariant.

Since Γ is quasi-isometric to G , the quasi-action of $\tilde{\Gamma}_0$ on G is proper, which means $\hat{\Gamma}_0$ and $\ker(h)$ quasi acts properly on $\prod_{[\Xi]} (\partial_{[\Xi]}^-, D_{[\Xi]})$. Now by Theorem 18 of [D], Γ is virtually polycyclic. $\square$

In a group G , an element $x \in G$ is called *exponentially distorted* if there are numbers c, ϵ such that for all $n \in \mathbb{Z}$,

$$\frac{1}{c} \log(|n| + 1) - \epsilon \leq \|x^n\|_G \leq c \log(|n| + 1) + \epsilon$$

where $\|x^n\|_G$ is the distance between the identity and x^n in G . In the case of a connected, simply connected solvable Lie group G , Osin showed in [O] that the set of exponentially distorted elements forms a normal subgroup $R_{\exp}(G)$.

Lemma 5.3.1. *Let G be a connected, simply connected solvable Lie group such that*

$$1 \rightarrow R_{\exp}(G) \rightarrow G \rightarrow \mathbb{R}^s \rightarrow 1,$$

where $R_{\exp}(G)$ is abelian. Then the above sequence splits and G is a semidirect product of $R_{\exp}(G)$ and $\mathbb{R}^s$.

Proof. Let $\mathfrak{h}$ be a Cartan subalgebra of $\mathfrak{g}$, the Lie algebra of G . Then $\mathfrak{v}$, the Lie algebra of $R_{\exp}(G)$, is generated by root spaces in the decomposition of $\mathfrak{g}$ with respect to $\mathfrak{h}$. Since this is abelian, it means that $\mathfrak{g}$ is a semidirect product of $\mathfrak{h}$ and $\mathfrak{v}$. Since $\mathfrak{g}/\mathfrak{v}$ is abelian, $\mathfrak{h}$ is abelian. $\square$

Corollary 5.3.3. *Let $G = \mathbf{H} \rtimes_\psi \mathbf{A}$ be a non-degenerate, unimodular, split abelian-by-abelian solvable Lie group where ψ is diagonalizable. If Γ is a finitely generated group quasi-isometric to G , then Γ is virtually a lattice in a unimodular semidirect product of $\mathbf{H}$ and $\mathbf{A}$.*

Proof. By Corollary 5.3.2, Γ contains a finite index subgroup that is polycyclic. By a theorem of Mostow (Theorem 4.28 in [R]) which says that a polycyclic group contains a finite index subgroup that embeds as a lattice in a connected, simply connected Lie group, we have $\mathcal{L}$, a connected, simply connected solvable Lie group to which Γ is virtually a lattice of. The crux of the proof consists of showing that $\mathcal{L}$ satisfies the short exact sequence in Lemma 5.3.1, and the argument is practically

²When rank of G is 2 or higher, $t_{[\Xi], \gamma}$ equals to $e^{\Xi_0(\mathbf{t}_{0, \gamma})}$; but in rank 1 all that we can say is that it is of the form $e^{\Xi_0(\mathbf{s}_\gamma)}$ where $\mathbf{s}_\gamma$ might not be the same as $\mathbf{t}_{0, \gamma}$

that of Section 4.3 in [D] with minor modifications. We reproduce the skeleton of the proof below, and refer the readers to the relevant sections in [D] for details.

We are now going to construct a continuous homomorphism $\tilde{h} : \mathcal{L} \rightarrow \mathbf{A} = \mathbb{R}^n$ that is onto, whose finitely generated kernel not only is quasi-isometric to $\mathbf{H}$ but also coincides with the exponential radical of $\mathcal{L}$. As any finitely generated in the same quasi-isometric class as $\mathbb{R}^n$ must be virtually $\mathbb{R}^n$, by Lemma 5.3.1, $\mathcal{L}$ is virtually a semidirect product of $\mathbf{A}$ and $\mathbf{H}$. If this semidirect product were not unimodular, then $\mathcal{L}$ would be non-amenable, which is a contradiction because amenability is preserved under quasi-isometry.

The $\tilde{h}$ is going to be the composition of the following three homomorphisms:

- A. $\mathcal{L} \rightarrow \prod_{[\alpha]} QSim(\partial_{[\alpha]}^-)$
- B. Conjugation of a uniform subgroup of $\prod_{[\alpha]} QSim(\partial_{[\alpha]}^-)$ into $AIsoM(G)$, where $AIsoM(G)$ is the set of all maps of the form $\psi_\gamma((\mathbf{x}_{[\Xi]}))_{[\Xi]}, \mathbf{t}) = ((g_{[\Xi], \gamma}(\mathbf{x}_{[\Xi]}))_{[\Xi]}, \mathbf{t} + \mathbf{s}_\gamma)$.
- C. $h : \prod_{[\alpha]} AIsoM(\partial_{[\alpha]}^-) \rightarrow \mathbf{A} = \mathbb{R}^n$

Homomorphism A. We can assume without loss of generality, that Γ itself is a lattice in $\mathcal{L}$. We start with the following construction which can be found in Section 3.2 of [F]. Choose some open subset $E \subset \mathcal{L}$ with compact closure, such that $\mathcal{L}$ is the union of left translates of E by Γ . Also fix a function $p : \mathcal{L} \rightarrow \Gamma$ such that $x \in p(x)E$ for every $x \in \mathcal{L}$. Then by defining $q_h : \Gamma \rightarrow \Gamma$ as $q_h(\gamma) = p(h\gamma)$ for every $h \in \mathcal{L}$, we obtain a homomorphism from $\mathcal{L}$ into $QI(\Gamma)$. Since Γ is quasi-isometric to G , conjugating by the quasi-isometry between Γ and G , we obtain a homomorphism from $\mathcal{L}$ into $QI(G)$, where the images have uniform quasi-isometric constants. By (v) of Proposition 5.3.1, we can realize $QI(G)$ as a subgroup of $\prod_{[\alpha]} QSim(\partial_{[\alpha]}^-)$. By passing to a finite index subgroup of $\mathcal{L}$ if necessary, we now have the homomorphism A from $\mathcal{L}$ to $\prod_{[\alpha]} QSim(\partial_{[\alpha]}^-)$, whose image is a uniform subgroup of quasi-similarities. Continuity of homomorphism A follows from Proposition 26 of [D] where continuity in each factor was obtained.

Homomorphism B. Theorem 2 of [D] says that we can conjugate the image of homomorphism A. into $\prod_{[\alpha]} ASim(\partial_{[\alpha]}^-)$. That elements of $\prod_{[\alpha]} ASim(\partial_{[\alpha]}^-)$ can be realized as elements of $AIsoM(G)$ follows from the **Claim** in the proof of Corollary 5.3.2. Homomorphism B. is continuous because conjugation is continuous.

Homomorphism C. The definition of $AIsoM(G)$ means that we have a well-defined homomorphism into $\mathbf{A} = \mathbb{R}^n$, which is our homomorphism C. Now if q_i is a sequence in $AIsoM(G)$ approaching to identity, then the map each one of them induces on the $\mathbf{A}$ factor also has to approach that of what the identity does. Since the identity map produces no change in the $\mathbf{A}$ factor, it follows that the image of q_i 's under homomorphism C. approaches $\vec{0} \in \mathbf{A}$.

Since Γ is quasi-isometric to G , the quasi-action³ of Γ , and therefore $\mathcal{L}$, on G is cobounded, it follows that $\tilde{h}$ must be onto because it is continuous.

We now claim that $R_{exp}(\mathcal{L}) = \ker(\tilde{h})$. To this end, we need the following from [O].

³Conjugating each left translation of Γ by the quasi-isometry between Γ and G gives a quasi-action on G

Lemma 5.3.2. (Lemma 2.1 in [O])

Suppose G, H are locally compact groups generated by some symmetric compact neighborhoods of the identities, $\|\cdot\|_G, \|\cdot\|_H$ are canonical norms on G and H , and $\text{dis}_G, \text{dist}_H$ are the induced metrics. Assume $\phi : G \rightarrow H$ is a continuous surjective homomorphism, then there is a constant K such that

$$\text{dist}_H(\phi(g_1), \phi(g_2)) \leq K \text{dis}_G(g_1, g_2)$$

Applying the lemma above to $\tilde{h}$ gives us that

$$\|\tilde{h}(\gamma)\| \leq K|\gamma|_{\mathcal{L}}, \forall \gamma \in \mathcal{L}$$

Now let $\gamma \in R_{\text{exp}}(\mathcal{L})$ such that $|\tilde{h}| = c$. Then for any n

$$cn = |\tilde{h}(\gamma^n)| \leq K|\gamma^n|_{\mathcal{L}} = K \log(n+1)$$

So we must have $\tilde{h} = \vec{0}$, hence $R_{\text{exp}}(\mathcal{L}) \subset \ker(\tilde{h})$.

Conulier showed in [C] that for a connected, simply connected solvable Lie group X , the *asymptotic dimension*, defined as the dimension of $X/R_{\text{exp}}(X)$ is a quasi-isometric invariant. This means that

$$\dim \mathcal{L}/R_{\text{exp}}(\mathcal{L}) = \dim G/R_{\text{exp}}(G) = \dim \mathbf{A} = n$$

However as $\tilde{h}$ is onto, the dimension of $\mathcal{L}/\ker(\tilde{h})$ also equals n . So $\ker(\tilde{h})$ cannot be strictly bigger than $R_{\text{exp}}(\mathcal{L})$.

By construction, $\mathcal{L}$ quasi-acts properly on G as a uniform group of quasi-isometries, which means $\ker(\tilde{h})$ quasi-acts properly on $\mathbf{H}$ as a uniform group of quasi-similarities, so $\ker(\tilde{h})$ is finitely generated by Proposition 20 in [D]. Fix a $p \in G$. Then $\gamma \mapsto B \circ A(\gamma)(p)$ is a quasi-isometry from $\mathcal{L}$ to G . Here B, A refers to the homomorphisms mentioned above. The restriction of this map to $\ker(\tilde{h}) = R_{\text{exp}}(\mathcal{L})$ produces a quasi-isometric embedding of $R_{\text{exp}}(\mathcal{L})$ into $\mathbf{H}$. However since the cohomological dimension is a quasi-isometry invariant [G], the dimension of $R_{\text{exp}}(\mathcal{L})$ must equal that of $\mathbf{H}$. Now by theorem 7.6 of [FM], this embedding must be coarsely onto, which means $R_{\text{exp}}(\mathcal{L})$ is quasi-isometric to $\mathbf{H}$, so $R_{\text{exp}}(\mathcal{L})$ must be virtually $\mathbf{H}$ since the latter is abelian. $\square$

Example of a unimodular solvable Lie group not Q.I. to any finitely generated groups

We need the following result which is stated in [EFW1], and whose proof is finished in [D].

Theorem 5.3. *Theorem 1 in [D] If the rank of G is 1, then a finitely generated group Γ quasi-isometric to G is virtually a lattice in G .*

Let G be a rank 1 group with weights $1, 1, -2$. It has no lattice⁴ so Theorem 5.3 says that it cannot be quasi-isometric to any finitely generated groups.

⁴because if an elements of $SL_3(\mathbb{Z})$ has two distinct eigenvalues, one of them repeated twice, then they have to be $-1, -1, 1$. See [H]. But the diagonal matrix with those eigenvalues as entries is not conjugate to the action of $\mathbb{R}$ on $\mathbb{R}^3$ i.e. the diagonal matrix with entries e^1, e^1, e^{-2} .

References

- [BH] M. Bridson, A. Haefliger. Metric Spaces of non-positive curvature. Springer-Verlag Berlin Heidelberg 1999
- [C] Y. de Cornulier. Dimension of asymptotic cones in Lie groups.
- [D] T. Dymarz. Large scale geometry of certain solvable groups. to appear in GAFA.
- [EFW0] A. Eskin, D. Fisher, K. Whyte. Quasi-isometries and rigidity of solvable groups. Preprint.
- [EFW1] A. Eskin, D. Fisher, K. Whyte. Coarse differentiation of quasi-isometries I: spaces not quasi-isometric to Cayley graphs. *Pur. Appl. Math. Q.*
- [EFW2] A. Eskin, D. Fisher, K. Whyte. Coarse differentiation of quasi-isometries II: Rigidity for Sol and Lamplighter groups, preprint
- [F] A. Furman. Mostow-Margulis rigidity with locally compact targets, *Geom. Funct. Anal. (GAFA)* 11 (2001), 30-59.
- [FM] B. Farb, L. Mosher. On the asymptotic geometry of abelian-by-cyclic groups. *Acta Math.* , Vol. 184, No.2 (2000), 145-202.
- [G] S.M. Gersten. Quasi-isometry invariance of cohomological dimension, *Comptes Rendues Acad.Sci. Paris. Serie 1 Math* 316 (1993), 411-416
- [H] K. Hasegawa. Four dimensional compact solvmanifold with and without complex analytic structures.
- [K] A.W.Knapp. Lie groups beyond an introduction. Birkhauser. 2002
- [R] M. Raghunathan. Discrete subgroups of Lie groups. Springer-Verlag 1979
- [O] D. Osin. Exponential radicals of solvable Lie groups, *J. Algebra* 248 (2002), 790-805.
- [P] I. Peng. Coarse differentiation and quasi-isometries of a class of solvable Lie groups I. Preprint.